

Abemaciclib (Verzenio)

About This Medicine

ABEMACICLIB (a BEM a SYE klib) treats breast cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Pain in your abdomen
- Tiredness
- Infection
- Decreased appetite (decreased hunger)
- Headache
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in 20% or greater of people treated with abemaciclib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe diarrhea
- Severe decrease in the number of white blood cells and neutropenic fever - a type of fever that can develop when you have a very low number of white blood cells which can be life-threatening.
- Severe changes in your liver function

- Blood clots including events such as stroke and heart attack, which may be life-threatening. A blood clot in your leg may cause your leg to swell, appear red and warm, and/or cause pain. A blood clot in your lungs may cause trouble breathing, pain when breathing, and/or chest pain.
- Inflammation (swelling) and/or scarring of the lungs, which may be life-threatening. You may have a cough and/or trouble breathing.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- It is important that you notify your care team at the first sign of diarrhea so they can provide you with anti-diarrheal medicine and give you further instructions. Notify your care team if you are taking anti-diarrheal medicine and your symptoms have not improved or are worsening after 24 hours.

How to Take Your Medicine

- Swallow the medicine whole with or without food. Take this medicine at the same time each day.
- Do not chew, crush or split tablets.
- If any of the tablets are broken, cracked or otherwise not intact, do not take them. Carefully throw away the tablets and wash your hands after handling.
- **Missed dose:** If you miss or vomit a dose, do not take the missed dose at all and do not double up on the next dose. Instead, continue with your regular dosing schedule.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.



- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop your diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your care team and they can help you get in touch with programs in your local area.

Food and Medicine Interactions

- Avoid grapefruit or grapefruit juice while taking this medicine. Grapefruit and grapefruit juice may raise the levels of abemaciclib in your body. This could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with abemaciclib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

When to Call Your Care Team



Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Headache that does not go away
- Easy bleeding or bruising
- Your leg is swollen, red, warm and/or painful
- Pain in your chest
- Dry cough
- Wheezing and/or trouble breathing
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Stroke. Symptoms of a stroke such as sudden numbness or weakness of your face, arm, or leg, especially on one side of your body; sudden confusion, trouble speaking or understanding; sudden trouble seeing in one or both eyes; sudden trouble walking, feeling dizzy, loss of balance or coordination; or sudden bad headache with no known cause. **If any of these symptoms last 2 minutes, call 911.**
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your abdomen that does not go away
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, trouble urinating, burning or pain when you pass urine, redness and/or swelling of the skin

Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 3 weeks after the last dose. Tell your care team right away if you think you might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for at least 3 weeks after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.



Revised December 2024

