

Adagrasib (Krazati)

About This Medicine

ADAGRASIB (a DA gra sib) treats lung cancer and colorectal cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Nausea and vomiting
- Diarrhea (loose bowel movements)
- Tiredness
- Swelling of the hands, feet, or any other part of the body
- Decreased appetite (decreased hunger)
- Muscle and bone pain
- Changes in your liver function
- Changes in your kidney function
- Trouble breathing

Note: Each of the side effects above was reported in 25% or greater of people treated with adagrasib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe nausea, vomiting and diarrhea, including colitis, which is inflammation (swelling) in the colon - symptoms are diarrhea (loose bowel movements), stomach cramping, and sometimes blood in the bowel movements. You may also be at risk of developing a partial or complete blockage of your small and/or large intestine.
- Abnormal electrocardiogram (EKG/ECG)
- Severe changes in your liver function

- Inflammation and/or scarring of the lungs, which can be life-threatening. You may have a cough and/or trouble breathing.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- It is important that you notify your care team at the first sign of diarrhea, nausea or vomiting so they can provide you with anti-diarrheal or anti-nausea medicine and give you further instructions. Notify your care team if you are taking anti-diarrhea medicine and your symptoms have not improved or are worsening.

How to Take Your Medicine

- Swallow the medicine whole with or without food. Do not chew, crush or split it. Take it at approximately the same time each day.
- **Missed dose:** If you miss a dose by less than 4 hours, take the missed dose right away and go back to your normal schedule. If you miss a dose by more than 4 hours, then skip the missed dose and go back to your normal schedule. Do not replace a vomited dose and take your next dose at the regular time.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Do not remove the desiccant (drying agent) from the container after opening. Do not eat or swallow the desiccant. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- Drink enough fluids to keep your urine pale yellow.



- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.

Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your care team as this could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with adagrasib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Pain in your chest
- Dry cough
- Wheezing and/or trouble breathing
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Tiredness and/or weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your abdomen that does not go away
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy

- Blood in your stool
- No bowel movement in 3 days or when you feel uncomfortable.
- Unable to pass gas
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of the hands, feet, or any other part of the body
- Pain that does not go away or is not relieved by prescribed medicine
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Decreased or very dark urine

Reproduction Warnings

- **Pregnancy warning:** Talk with your care team if you are pregnant or planning to become pregnant while taking this medicine. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 week after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised February 2025

