

## Aldesleukin (Proleukin)

---

### About This Medicine

ALDESLEUKIN (al des LOO kin) treats kidney cancer and skin cancer. It works by helping your immune system slow or stop the spread of cancer cells. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

### Possible Side Effects

- Decrease in the number of platelets. This may raise your risk of bleeding.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Chills
- Changes in your liver function
- Confusion
- Changes in your kidney function and/or decreased urine
- Trouble breathing
- Rash
- Low blood pressure

**Note:** Each of the side effects above was reported in 30% or greater of people treated with aldesleukin. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

### Warnings and Precautions

- A life-threatening syndrome where fluid and protein can leak from your blood vessels into your tissues. This can cause a decrease in your blood protein level and blood pressure and fluid can accumulate in your tissues and/or lungs.
- Changes in your central nervous system can happen. The central nervous system is made up of your brain and spinal cord. You could feel extreme tiredness, agitation, confusion, hallucinations (see or hear things that are not there), trouble understanding or speaking, loss of control of your bowels or bladder, eyesight changes, numbness, or lack of strength to your arms, legs, face, or

body, and coma. **If you start to have any of these symptoms let your care team know right away.**

- Changes in your white blood cells that may increase your risk of severe and life-threatening infections
- If you have an autoimmune disease or disorder that causes inflammation, it may get worse with aldesleukin.
- Blood sugar levels may change, and you may develop diabetes. If you have diabetes, changes may need to be made to your diabetes medicine.
- Changes in your thyroid function
- Changes in your heart function such as inflammation (swelling) in the tissue of the heart, heart attack and/or congestive heart failure – your heart is not pumping blood as well as it should be, and fluid can build up in your body.
- Allergic reactions, including anaphylaxis, are rare but may happen in some people. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**
- **Infusion-related reaction:** While you are getting this medicine in your vein (IV), you may have a reaction to the medicine. Sometimes you may be given medicine to stop or lessen these side effects. Your care team will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**

**Note:** *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.*

## Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- Taking this medicine can cause an increased risk of having an allergic reaction to iodine contrast material. This may happen while you are receiving treatment with aldesleukin and for several months after you get the last dose of this medicine. Before you have a procedure with an iodine contrast material, talk with your care team for further instructions.

## Treating Side Effects

- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever



- To decrease your risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have loose bowel movements, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop your diarrhea.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- If you have diabetes, keep good control of your blood sugar level. Tell your care team if your glucose levels are higher or lower than normal.
- Infusion reactions may occur after your infusion. **If this happens, call 911 for emergency care.**

## Food and Medicine Interactions

- There are no known interactions of aldesleukin with food.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with aldesleukin. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Easy bleeding or bruising
- Feeling dizzy or lightheaded
- Tiredness that interferes with your daily activities
- Confusion and/or agitation
- Hallucinations
- Symptoms of a seizure such as confusion, blacking out, passing out, loss of hearing or vision, blurred vision, unusual smells or tastes (such as burning rubber), trouble talking, tremors or shaking in parts or all of the body, repeated body movements, tense muscles that do not relax, and loss of control of urine and bowels. **If you or your family member suspects you are having a seizure, call 911 right away.**



- Trouble understanding or speaking
- Blurry vision or changes in your eyesight
- Numbness or lack of strength to your arms, legs, face, or body
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Wheezing and/or trouble breathing
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back or it can be constant. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Cough that is bothersome
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Decreased and/or very dark urine
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of the hands, feet, or any other part of the body
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- New rash and/or a rash that is not relieved by prescribed medicines
- Any change in a pre-existing autoimmune or inflammatory condition
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**
- Signs of low thyroid function: tiredness, unexplained weight gain, hair loss, dry skin, constipation, increased sensitivity to cold
- Signs of increased thyroid function: unexplained weight loss, sweating, irritability, feeling that your heart is beating fast or in a not normal way (palpitations)
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, trouble urinating, burning or pain when you pass urine, redness and/or swelling of the skin

## Reproduction Warnings

- **Pregnancy warning:** Talk with your care team if you are pregnant or planning to become pregnant while taking this medicine. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Talk with your care team about breastfeeding during treatment. You may need to stop breastfeeding.



- **Fertility warning:** The effect of this medicine on fertility is not known. If you plan to have children, talk with your care team.

Revised January 2026

