

## Anti-thymocyte globulin (rabbit) (Thymoglobulin)

---

### About This Medicine

ANTITHYMOCYTE IMMUNE GLOBULIN (an tee THI mo cite im MUNE GLOB yoo lin) prevents the body from rejecting an organ transplant. It works by lowering the body's immune system response. This helps the body accept the donor organ. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

### Possible Side Effects

- Decrease in the number of white blood cells and platelets. This may raise your risk of infection and your risk of bleeding.
- Nausea
- Pain in your abdomen
- Fever and chills
- Increased potassium in your blood
- Headache
- Feeling nervous or worried (anxiety)
- Urinary tract infection
- Trouble breathing
- High blood pressure

**Note:** Each of the side effects above was reported in greater than 5% of people treated with anti-thymocyte globulin (rabbit). All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

### Warnings and Precautions

- Allergic reactions, including anaphylaxis are rare but may happen in some people and can be life-threatening. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**
- This medicine may raise your risk of getting another cancer, such as lymphoma.



- Severe decrease in white blood cells and/or platelets.
- Risk of severe and life-threatening infections.
- Injection site reaction - you may get a rash, swelling or bruising or your skin may get red, warm, itchy, or painful at the site of your infusion.
- While you are getting this medicine in your vein (IV), you may have a reaction to the medicine. Sometimes you may be given medicine to stop or lessen these side effects. Your care team will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Cytokine release syndrome (CRS), which can be life-threatening. Some types of cancer drugs can cause CRS because of the effects of the drug in your body. If this happens you may feel very sick and get a fever, headache, nausea, or feel weak. You may also have changes to your blood pressure. Because of this, your blood pressure and pulse may be checked while you are getting this drug.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

## Important Information

- Talk to your care team before receiving any vaccinations during your treatment. Some vaccinations are not recommended while receiving anti-thymocyte globulin (rabbit).

## Treating Side Effects

- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- Drink enough fluids to keep your urine pale yellow.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- Cytokine release syndrome may happen after your infusion. **If this happens, call 911 for emergency care**
- Infusion reactions may occur after your infusion. **If this happens, call 911 for emergency care.**
- While you are getting this drug, please tell your care team right away if you get a rash, swelling or bruising or your skin gets red, warm, itchy or painful at the site of your infusion.



- Use sunscreen with SPF 30 or higher when you are outdoors even for a short time. Cover up when you are out in the sun. Wear wide-brimmed hats, long-sleeved shirts, and pants. Keep your neck, chest, and back covered.

## Food and Medicine Interactions

- There are no known interactions of anti-thymocyte globulin (rabbit) with food.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with anti-thymocyte globulin (rabbit). Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Blurry vision or other changes in eyesight
- Headache that does not go away
- Easy bleeding or bruising
- Wheezing and/or trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your abdomen that does not go away
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, trouble urinating, burning or pain when you pass urine, redness and/or swelling of the skin
- Signs of a urinary tract infection such as burning or pain when you pass urine, trouble urinating, smelly and/or cloudy urine, pain in your lower abdomen or lower/side of your back.
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**
- Signs of cytokine release syndrome such as: fever, chills, or shaking chills, feeling dizzy or lightheaded, have a headache and/or have nausea or throwing up. **If this happens, call 911 for emergency care.**
- Signs of high potassium: muscle weakness, nausea, vomiting, diarrhea, chest pain and/or feeling that your heart is beating in a fast or not normal way.



## Reproduction Warnings

- **Pregnancy warning:** Talk with your care team if you are pregnant or planning to become pregnant while taking this medicine. If you can become pregnant, use birth control while taking this medicine and for at least 3 months after the last dose. Tell your care team right away if you think you might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine.
- **Fertility warning:** The effect of this medicine on fertility is not known. If you plan to have children, talk with your care team.

Revised December 2025

