

Asparaginase Erwinia chrysanthemi (recomb)-rywn (Rylaze)

About This Medicine

ASPARAGINASE ERWINIA CHRYSANTHEMI (AS par a JIN ase er WIN i a kris an THEM i) treats leukemia and lymphoma. It works by slowing down the growth of cancer cells. It is given as an injection in your muscle (intramuscularly).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea
- Diarrhea (loose bowel movements)
- Pain in your abdomen
- Decreased appetite (decreased hunger)
- Tiredness
- Infection
- Fever and neutropenic fever - a type of fever that can develop when you have a very low number of white blood cells which can be life-threatening.
- Allergic reactions, including anaphylaxis, are rare but may happen in some people. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**
- Changes in your liver function
- Inflammation (swelling) of your pancreas
- Decreased level of potassium in your blood
- Blood sugar levels may change
- Abnormal bleeding – symptoms may be coughing up blood, throwing up blood (may look like coffee grounds), red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding.
- Muscle and bone pain
- Headache

Note: Each of the side effects above was reported in greater than 20% of people treated with asparaginase erwinia chrysanthemi (recomb)-rywn. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe allergic reactions
- Severe inflammation of your pancreas, which can be life-threatening
- Severe changes in your liver function and blockage of small veins in the liver, which can cause liver failure and be life-threatening.
- Blood clots and events such as stroke and heart attack. A blood clot in your leg may cause your leg to swell, appear red and warm, and/or cause pain. A blood clot in your lungs may cause trouble breathing, pain when breathing, and/or chest pain.
- Severe abnormal bleeding

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- Drink enough fluids to keep your urine pale yellow.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.



- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- If you have diabetes, keep good control of your blood sugar level. Tell your care team if your glucose levels are higher or lower than normal.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.

Food and Medicine Interactions

- There are no known interactions of asparaginase erwinia chrysanthemi (recomb)-rywn with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine interacts with hormonal birth control (i.e., birth control pills, skin patches, vaginal rings, injections) and affect how they work. Discuss with your care team what method of birth control may be right for you during your treatment.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Headache that does not go away
- Wheezing and/or trouble breathing
- Tiredness and/or extreme weakness that interferes with normal activities
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Symptoms of a stroke such as sudden numbness or weakness of your face, arm, or leg, especially on one side of your body; sudden confusion, trouble speaking or understanding; sudden trouble seeing in one or both eyes; sudden trouble walking, feeling dizzy, loss of balance or coordination; or



sudden bad headache with no known cause. **If you have any of these symptoms for 2 minutes, call 911.**

- Pain in your mouth or throat that makes it hard to eat or drink
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Severe pain in your abdomen that does not go away and that may spread to your back
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, trouble urinating, burning or pain when you pass urine, redness and/or swelling of the skin
- Signs of abnormal bleeding such as coughing up blood, throwing up blood (may look like coffee grounds), easy bleeding or bruising, red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Your leg is swollen, red, warm, and/or painful
- Pain that does not go away, or is not relieved by prescribed medicines

Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use non-hormonal birth control while taking this medicine and for 3 months after the last dose. Tell your care team right away if you think you might be pregnant.
- Birth control pills (oral contraceptives) and other hormonal forms of birth control may not be effective with this medicine.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 week after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised August 2025

