

Belantamab mafodotin-blmf (Blenrep)

About This Medicine

BELANTAMAB MAFODOTIN (bel AN ta mab MA foe DOE tin) treats multiple myeloma, a type of bone marrow cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Changes to the cornea (seen on eye exam)
- Cataract (clouding of the lens in your eye)
- Eye irritation, feeling like something is in your eye
- Blurred or decreased vision
- Dry and/or painful eyes
- Sensitivity to light (photosensitivity). Photosensitivity means that you may become more sensitive to the sun and/or light. Your eyes may water more, mostly in bright light.
- Diarrhea (loose bowel movement)
- Tiredness
- Increase in creatine phosphokinase (CPK), which can sometimes be the result of muscle inflammation or damage. You may have muscle aching and/or cramping.
- Changes in your liver and/or kidney function
- Upper respiratory tract infection
- Pneumonia
- COVID-19

Note: Each of the side effects above was reported in greater than 20% of people treated with belantamab mafodotin-blmf. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe vision problems such as dry eyes, blurred vision, worsening vision and/or vision loss, and ulcers on your cornea.
- Decrease in the number of platelets. This may raise your risk of bleeding and can be life-threatening.

Note: *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.*

Important Information

- This medicine is part of special program which helps make sure it is used safely (Risk Evaluation and Mitigation Strategy [REMS] program). As part of this program, you may be asked to read safety information or complete certain tests while taking it. Your care team will provide you with more information.
- This medicine may impair your ability to drive or use machinery. Use caution and talk to your care team about any precautions you may need to take.
- You should use preservative-free lubricant eye drops at least 4 times a day during your treatment. Ask your care team for more information about which product they recommend for you.
- It is recommended to avoid wearing contact lenses during treatment unless told otherwise by your eye specialist.
- It is important that you see an eye specialist before and during your treatment, even if your vision seems fine. Discuss with your care team or more information.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Wear dark sunglasses when in the sun or bright lights
- Drink enough fluids to keep your urine pale yellow.



- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop diarrhea.

Food and Medicine Interactions

- There are no known interactions of belantamab mafodotin-blmf with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Feeling like something is in your eyes or eye irritation
- Blurred vision or any other changes in eyesight
- Sensitivity to light
- Dry or painful eyes
- Wheezing and/or trouble breathing
- Coughing that is bothersome or coughing up yellow, green, or bloody mucus
- New loss of taste or smell
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Decreased or very dark urine
- Unexplained muscle pain, tenderness, or weakness

Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 4 months after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 6 months after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.



- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 3 months after the last dose.

Fertility warning: This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised November 2025

