

Capmatinib (Tabrecta)

About This Medicine

CAPMATINIB (kap MA ti nib) treats lung cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team you have questions.

Possible Side Effects

- Nausea and vomiting (throwing up)
- Tiredness
- Swelling of the hands, feet, or any other part of the body
- Decreased appetite (decreased hunger)
- Bone and muscle pain
- Cough
- Trouble breathing

Note: Each of the side effects above was reported in 20% or greater of people treated with capmatinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Inflammation (swelling) and/or scarring of the lungs, which can be life-threatening. You may have a cough and/or trouble breathing.
- Changes in your pancreas function and/or inflammation of your pancreas
- Changes in your liver function
- Sensitivity to sunlight/light. You may get a skin rash/reaction if you are in the sun or are exposed to sun lamps and tanning beds.
- Allergic reactions are rare but may happen in some people. Signs of allergic reaction to this medicine may be nausea, vomiting, rash, itching, fever, chills and feeling dizzy. **If this happens, do not take another dose of this drug. You should get urgent medical treatment.**

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

How to Take Your Medicine

- Swallow the medicine whole with or without food. Do not break, chew, or crush tablets.
- **Missed dose:** If you vomit or miss a dose, take your next dose at the regular time. Do not take 2 doses at the same time and do not double up on the next dose.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Protect from moisture and do not remove the desiccant. Throw away any unused tablets after 6 weeks of first opening the container. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Drink enough fluids to keep your urine pale yellow.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you throw up, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- Use sunscreen with SPF 30 or higher when you are outdoors even for a short time. Cover up when you are out in the sun. Wear sunglasses, wide-brimmed hats, long-sleeved shirts, and pants. Keep your neck, chest, and back covered.



- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.

Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your care team as this could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with capmatinib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Wheezing and/or trouble breathing
- Pain in your chest
- Dry cough and/or a cough that is bothersome
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Severe pain in your abdomen that may spread to your back
- Pain that does not go away, or is not relieved by prescribed medicines
- Extreme weakness that interferes with normal activities
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of the hands, feet, or any other part of the body
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- A new rash or a rash that is not relieved by prescribed medicines
- Develop sensitivity to sunlight/light

Reproduction Warnings



- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 1 week after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 1 weeks after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 week after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised December 2025

