

Cladribine

About This Medicine

CLADRIBINE (KLA dri been) treats leukemia. It works by slowing down the growth of cancer cells. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Nausea
- Tiredness
- Fever
- Infection
- Headache
- Rash
- Injection site reaction you may get a rash, swelling or bruising or your skin may get red, warm, itchy, or painful at the site of your infusion.

Note: All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe muscle weakness in your arms and/or legs which can be irreversible.
- Effects on the nerves are called peripheral neuropathy. You may feel numbness, tingling, or pain
 in your hands and feet. It may be hard for you to button your clothes, open jars, or walk as usual.
 The effect on the nerves may get worse with more doses of the medicine. These effects get better
 in some people after the medicine is stopped but it does not get better in all people.
- Severe bone marrow suppression and neutropenic fever, a type of fever that can develop when you
 have a very low number of white blood cells which can be life-threatening.
- · Risk of severe and life-threatening infections



Cladribine continued Page 2 of 3

Tumor lysis syndrome: This medicine may act on the cancer cells very quickly. This may affect how
your kidneys work.

Changes in your kidney function

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- Talk to your care team before receiving any vaccinations during your treatment. Some vaccinations
 are not recommended while receiving cladribine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.
- While you are getting this medicine, please tell your care team right away if you get a rash, swelling or bruising or if your skin gets red, warm, itchy or painful at the site of the IV infusion.
- Keeping your pain under control is important to your well-being. Please tell your care team if you
 are experiencing pain.



Cladribine continued Page 3 of 3

Food and Medicine Interactions

- There are no known interactions of cladribine with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- · Tiredness that interferes with your daily activities
- · Feeling dizzy or lightheaded
- Headache that does not go away
- Weakness or lack of strength to your arms and/or legs
- Easy bleeding or bruising
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- · Decreased or very dark urine
- Signs of tumor lysis: confusion or agitation, decreased urine, nausea/vomiting, diarrhea, muscle cramping, numbness and/or tingling, seizures.
- Numbness, tingling, or pain in your hands and feet
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, difficulty urinating, burning or pain when you pass urine, redness and/or swelling of the skin
- A new rash and/or itching or a rash that is not relieved by prescribed medicines

Reproduction Warnings

- **Pregnancy warning**: This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine. Let your care team right away if you think you might be pregnant.
- Breastfeeding warning: Talk with your care team about breastfeeding during treatment. You may need to stop breastfeeding.
- **Fertility warning:** The effect of this medicine on fertility is not known. If you plan to have children, talk with you care team.

Revised July 2025

