

## Daratumumab and hyaluronidase-fihj (Darzalex Faspro)

---

### About This Drug

Daratumumab and hyaluronidase-fihj is used to treat cancer. It is given as an injection under the skin (subcutaneously).

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Upper respiratory tract infection

**Note:** Each of the side effects above was reported in 20% or greater of patients treated with daratumumab and hyaluronidase-fihj. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

### Warnings and Precautions

- Severe decrease in the number of white blood cells and platelets.
- While you are getting this injection, you may have a reaction to the drug, which may be life-threatening. Sometimes you may be given medication to stop or lessen these side effects. Your nurse will check you closely for these signs: fever or shaking chills, flushing, facial swelling, blurred vision or other changes in eyesight, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Injection site reaction - you may get a rash, swelling, or bruising or your skin may get red, warm, itchy, or painful at the site of your injection.
- Severe and life-threatening heart problems in patients receiving daratumumab and hyaluronidase-fihj with light chain (AL) amyloidosis
- This medication can affect the results of blood tests that match your blood type during your treatment and for up to 6 months after your treatment. Your blood type will be tested before treatment. Be sure to tell all healthcare providers you are taking this medicine before receiving blood transfusions, and for 6 months after your last dose.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

## Important Information

- This drug may reactivate the hepatitis B virus. Talk to your doctor if you have ever had or might have a hepatitis B infection.
- If you are taking this drug in combination with lenalidomide, thalidomide, or pomalidomide, talk to your doctor and/or nurse about precautions you may need to take with blood and/or sperm donation.
- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- Injection-related reactions may rarely occur after your injection. **If this happens, call 911 for emergency care.**
- While you are getting this drug, please tell your nurse right away if you get a rash, swelling, or bruising or your skin gets red, warm, itchy, or painful at the site of your injection.

## Food and Drug Interactions

- There are no known interactions of daratumumab and hyaluronidase-fihj with food.
- This drug may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call the Doctor



Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Tiredness that interferes with your daily activities
- Coughing up yellow, green, or bloody mucus
- Wheezing and/or trouble breathing
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back, or it can be constant. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Signs of injection-related drug reaction: fever or shaking chills, flushing, facial swelling, blurred vision or other changes in eyesight, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**
- If you think you may be pregnant

## Reproduction Warnings

- **Pregnancy warning:** This drug may have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 3 months after stopping treatment. Let your doctor know right away if you think you may be pregnant. When receiving daratumumab and hyaluronidase-fihj in combination with other agents, please refer to the medication guide of those agents for more specific information.
- **Breastfeeding warning:** It is not known if this drug passes into breast milk. For this reason, women should talk to their doctor about the risks and benefits of breastfeeding during treatment with this drug because this drug may enter the breast milk and cause harm to a breastfeeding baby. When receiving daratumumab and hyaluronidase-fihj in combination with other agents, please refer to the medication guide of those agents for more specific information.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised July 2023

