

## Duvelisib (Copiktra)

---

### About This Medicine

DUVELISIB (DOO ve LIS ib) treats leukemia and lymphoma. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions.

### Possible Side Effects

- Decrease in the number of white blood cells and red blood cells. This may raise your risk of infection and make you tired and weak.
- Nausea
- Diarrhea (loose bowel movements)
- Colitis, inflammation (swelling) in the colon - symptoms are diarrhea, stomach cramping, and sometimes blood in the bowel movements
- Tiredness
- Fever
- Muscle and bone pain
- Cough
- Upper respiratory tract infection
- Pneumonia
- Rash

**Note:** Each of the side effects above was reported in 20% or greater of people treated with duvelisib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team

### Warnings and Precautions

- Risk of severe and life-threatening infections
- Severe diarrhea and colitis, which can be life-threatening
- Severe allergic skin reactions, which can be life-threatening. You may develop blisters on your skin that are filled with fluid or a severe red rash all over your body that may be painful and sometimes



peel off. Sometimes, you can have a rash, along with problems with your organs, swollen lymph nodes, fever and higher than normal white blood cells.

- Inflammation of the lungs, which can be life-threatening. You may have a dry cough or trouble breathing.
- Severe changes in your liver function
- Severe decrease in the number of white blood cells

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

## How to Take Your Medicine

- Swallow the medicine whole, with or without food. Do not open, break or chew the capsules.
- **Missed dose:** If you miss a dose by less than 6 hours, take it as soon as you think about it, and then take your next dose at the regular time. If you miss a dose by more than 6 hours, then skip the missed dose and take your next dose at the regular time.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- Drink enough fluids to keep your urine pale yellow.



- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that is available to help stop or lessen diarrhea.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.

## Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your care team as this could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs and others) you are taking before starting this medicine as there are known medicine interactions with duvelisib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

## When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and/or weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Wheezing and/or trouble breathing
- Dry cough and/or a cough that is bothersome
- Coughing up yellow, green, or bloody mucus
- Pain in your chest
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Pain in your abdomen that does not go away
- Blood in your stool (bright red, or black/tarry)
- New rash and/or itching



- Rash that is not relieved by prescribed medicines
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, trouble urinating, burning or pain when you pass urine, redness and/or swelling of the skin
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Flu-like symptoms: fever, headache, muscle and joint aches, and fatigue (low energy, feeling weak)
- Swollen lymph nodes in your neck and/or armpits
- Pain that does not go away, or is not relieved by prescribed medicines

## Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 1 month after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 1 month after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 month after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised March 2026

