

Enfortumab vedotin-ejfv (Padcev)

About This Drug

Enfortumab vedotin-ejfv is used to treat cancer. It is given in the vein (IV).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Tiredness
- Nausea
- Diarrhea (loose bowel movements)
- Changes in the way food and drinks taste
- Decreased appetite (decreased hunger)
- Weight loss
- Changes in your liver and kidney function
- Changes in your electrolytes
- Changes in your pancreas function
- Decrease in a blood protein called albumin
- An increased in uric acid in the blood
- Blood sugar level may change
- Effects on the nerves called peripheral neuropathy. You may feel numbness, tingling, or pain in your hands and feet. It may be hard for you to button your clothes, open jars, or walk as usual. The effect on the nerves may get worse with more doses of the drug. These effects get better in some people after the drug is stopped but it does not get better in all people.
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.
- Dry skin
- Itching
- Rash

Note: Each of the side effects above was reported in 20% or greater of patients treated with enfortumab vedotin-ejfv. All possible side effects are not included. Your side effects may be different depending on

your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Severe changes in your blood sugar level which may be life-threatening
- Severe effects on the nerves called peripheral neuropathy
- Changes in eyesight such as dry eyes and blurred vision
- Severe allergic skin reactions which can be life-threatening. You may develop blisters on your skin that are filled with fluid or a severe red rash all over your body that may be painful.
- Hand-foot syndrome. The palms of your hands or soles of your feet may tingle, become numb, painful, swollen, or red.
- Skin and tissue irritation including redness, pain, warmth, or swelling at the IV site if the drug leaks out of the vein and into nearby tissue.
- Inflammation (swelling) and/or scarring of the lungs, which can be life-threatening. You may have a dry cough or trouble breathing.

Note: *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.*

Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).



- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- Taking good care of your mouth may help food taste better and improve your appetite.
- To help with weight loss, drink fluids that contribute calories (whole milk, juice, soft drinks, sweetened beverages, milkshakes, and nutritional supplements) instead of water.
- If you have diabetes, keep good control of your blood sugar level. Tell your nurse or your doctor if your glucose levels are higher or lower than normal.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.
- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.
- Moisturize your skin several times a day.
- Avoid sun exposure and apply sunscreen routinely when outdoors.
- If you have dry eyes, you should use artificial tears, hydrating or lubricating eye gels to help prevent or treat dry eyes. Ask your nurse or doctor for more information about which product they recommend for you.

Food and Drug Interactions

- This drug may interact with grapefruit and grapefruit juice. Talk to your doctor as this could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with enfortumab vedotin-ejfv. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor



Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and/or weakness that interferes with your daily activities
- Easy bleeding or bruising
- Feeling dizzy or lightheaded
- Dry eye
- Blurred vision or other changes in eyesight
- Pain in your chest
- Dry cough
- Wheezing and/or trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- New rash and/or itching that is bothersome
- Dry skin that is bothersome
- Rash that is not relieved by prescribed medicines
- Numbness, tingling, or pain in your hands and/or feet
- Painful, red, or swollen areas on your hands or feet
- Swelling of the hands, feet, or any other part of the body
- Flu-like symptoms: fever, headache, muscle and joint aches, and fatigue (low energy, feeling weak)
- While you are getting this drug, please tell your nurse right away if you have any pain, redness, or swelling at the site of the IV infusion.
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Decreased or very dark urine
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 2 months after stopping treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 4 months after stopping treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for at least 3 weeks after stopping treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.



- **Fertility warning:** In men and women both, this drug may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised May 2023

