

Inavolisib (Itovebi)

About This Medicine

Inavolisib is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea
- Diarrhea (loose bowel movements)
- Tiredness
- Decreased appetite (decreased hunger)
- Blood sugar levels may change
- Electrolyte changes
- Changes in your kidney function
- Changes in your liver function
- COVID-19 infection
- Headache
- Rash

Note: Each of the side effects above was reported in 20% or greater of people treated with inavolisib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe changes in your blood sugar
- Severe soreness of the mouth and throat
- Severe diarrhea that can cause dehydration (lack of water in the body) and changes in your kidney function

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- It is important that you notify your care team at the first sign of diarrhea so they can provide you with anti-diarrheal medicine and give you further instructions. Notify your care team if you are taking anti-diarrhea medicine and your symptoms have not improved or are worsening.

How to Take Your Medicine

- Swallow the medicine whole with or without food. Take it at approximately the same time each day. Do not chew, split, or crush it.
- **Missed dose:** If you miss a dose by less than 9 hours, take the missed dose right away and go back to your normal schedule. If you miss a dose by more than 9 hours, then skip the missed dose and go back to your normal schedule. Do not replace a vomited dose.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.



- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day.
- Choose food and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping. Ask your care team about medicine that can lessen or stop diarrhea.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- If you have diabetes, keep good control of your blood sugar level. Tell your care team if your glucose levels are higher or lower than normal.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.

Food and Medicine Interactions

- There are no known interactions of inavolisib with food.
- Tell your care team about all the medicines and dietary supplements (vitamins, minerals, herbs and others) that you are taking at this time. The safety and use of dietary supplements and alternative diets are often not known. Using these might affect your cancer or interfere with your treatment. Until more is known, you should not use dietary supplements or alternative diets without your cancer care team's help.
- This medicine interacts with hormonal birth control (i.e., birth control pills, skin patches, vaginal rings, injections) and affect how they work. Discuss with your care team what method of birth control may be right for you during your treatment.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Headache that does not go away



- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Cough that is bothersome
- Trouble breathing
- New loss of taste or smell
- Pain in your mouth or throat that makes it hard to eat or drink
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Decreased or very dark urine
- Rash that is not relieved by prescribed medicines

Reproduction Warnings

- **Pregnancy warning:** This medicine may cause serious birth defects. If you can become pregnant, use non-hormonal birth control while taking this medicine and for 1 week after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 1 week after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- You should also refer to the prescribing information of each medicine you are also receiving for more specific information.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 week after the last dose. You should also refer to the prescribing information of each medicine you are also receiving for more specific information.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team. You should also refer to the prescribing information of each medicine you are also receiving for more specific information.

New October 2024

