

Lomustine (Gleostine)

About This Medicine

Lomustine is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea and vomiting (throwing up)
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Severe bone marrow suppression, which can be life-threatening.
- Thickening and/or inflammation (swelling) of the lung tissues. You may have a dry cough and/or trouble breathing.
- This medicine may raise your risk of getting a second cancer, such as acute leukemia and myelodysplasia.
- Changes in your liver function
- Changes in your kidney function, which can cause kidney failure.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

How to Take Your Medicine

- Swallow the medicine whole. Do not break the capsules.



- This medicine is usually taken a single dose that will not be repeated for at least 6 weeks. Each dose may be from different strengths and colors of capsules.
- **Missed dose:** If you vomit or miss a dose, contact your doctor for further instructions. Do not take 2 doses at the same time or extra doses.
- **Handling:** Wash your hands with soap and water before and after handling your medicine, yourself and your caretakers should not handle your medicine and your medicine bottle with bare hands and should wear latex gloves.
- If you get any of the contents of a broken capsules on your skin, you should wash the area of the skin well with soap and water right away. Call your doctor if you get a skin reaction.
- Cytotoxic medicines leave the body through urine and stool, but they can also be present in other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your health care provider after your treatment. If you take a cytotoxic pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Avoid temperatures over 40°C (104°F).
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.



- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.

Food and Medicine Interactions

- There are no known interactions of lomustine with food.
- This medicine may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Pain in your chest
- Dry cough
- Wheezing and/or trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your mouth or throat that makes it hard to eat or drink
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Decreased or very dark urine
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This medicine may have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 2 weeks after stopping treatment. Men with female partners of childbearing potential should use a condom during your cancer treatment and for 3.5 months after stopping treatment.



Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.

- **Breastfeeding warning:** Women should not breastfeed during treatment and for 2 weeks after stopping treatment because this medicine could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** In men and women both, this medicine may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised August 2023

