

Lusutrombopag (Mulpleta)

About This Medicine

Lusutrombopag is used to increase platelets. It is given orally (by mouth).

Possible Side Effects

- Headache

Note: The side effect above was reported in 3% or greater of patients treated with lusutrombopag. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Increased risk of blood clots, including blood clots in the liver. The risk may be increased if you have other known risks for blood clotting disorders.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

How to Take Your Medicine

- Swallow the medicine whole with or without food.
- **Missed dose:** If you miss a dose, take it as soon as you think about it on the day of the missed dose, otherwise skip the missed dose and take your next dose at the regular time. Do not take 2 doses at the same time and do not double up on the next dose.
- **Storage:** Store this medicine in the original container at room temperature.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects



- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.

Food and Medicine Interactions

- There are no known interactions of lusutrombopag with food.
- This medicine may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Headache that does not go away
- Wheezing and/or trouble breathing
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Symptoms of a stroke such as sudden numbness or weakness of your face, arm, or leg, mostly on one side of your body; sudden confusion, trouble speaking or understanding; sudden trouble seeing in one or both eyes; sudden trouble walking, feeling dizzy, loss of balance or coordination; or sudden, bad headache with no known cause. **If you have any of these symptoms for 2 minutes, call 911.**
- Symptoms of a clot in your liver such as pain in your abdomen, nausea and vomiting (throwing up), diarrhea (loose bowel movements), blood in your stools (bright red, or black/tarry).
- Your leg is swollen, red, warm and/or painful
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** This medicine may have harmful effects on the unborn baby. For this reason, be sure to talk with your doctor if you are pregnant or planning to become pregnant while receiving this medicine. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for at least 28 days after stopping treatment because this medicine could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this medicine. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.



Revised October 2023



User/Authorized User acknowledges that the ClinicalPath Portal is intended to be utilized as an information management tool only, and that Elsevier has not represented the ClinicalPath Portal as having the ability to diagnose disease, prescribe treatment, or perform any other tasks that constitute the practice of providing medical care. The clinical information contained within the ClinicalPath Portal is intended as a supplement to, and not a substitute for, the knowledge, expertise, skill, and judgment of physicians, pharmacists and other healthcare professionals involved with patient care at User/Authorized User facilities.
CONFIDENTIAL AND PROPRIETARY. NOT FOR DISTRIBUTION.