

## Lutetium Lu 177 dotatate (Lutathera)

---

### About This Medicine

LUTETIUM LU 177 DOTATATE (loo TEE shee uhm loo 177 DOE ta tate) treats neuroendocrine tumors, a cancer of the cells that release hormones and other substances in your body. It works by releasing energy (radiation) to slow or stop the spread of cancer cells. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Tiredness
- Nausea and vomiting (throwing up)
- Decreased appetite (decreased hunger)
- Pain in your abdomen
- Diarrhea (loose bowel movements)
- Changes in your liver function
- Changes in your kidney function
- Electrolyte changes
- Blood sugar levels may change
- Increased uric acid in the blood

**Note:** Each of the side effects above was reported in 20% or greater of people treated with lutetium Lu 177 dotatate. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

### Warnings and Precautions

- This medicine increases your exposure to radiation. Over time, long-term radiation exposure can increase your risk of getting another cancer. This medicine can also increase your risk of getting myelodysplastic syndrome and acute leukemia.
- Severe bone marrow suppression



- Allergic reactions, including anaphylaxis, are rare but may happen in some people. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, contact your care team and get urgent medical treatment.**
- Severe changes in your kidney function which can cause kidney failure
- Severe changes in your liver function which can cause liver failure
- While you are getting this medicine in your vein (IV), you may have a release of tumor-related hormones. You may experience sudden skin flushing, diarrhea, trouble breathing and low blood pressure. This reaction usually happens within the first 24 hours of your infusion.

**Note:** *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.*

## Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions and good hygiene practices during your treatment and for up to 30 days after treatment to minimize radiation exposure from your bodily fluids. Talk to your care team about the necessary precautions to take during this time.
- Drink plenty of fluids and try to urinate as often as possible to flush your bladder the day before treatment, on the day of treatment and the day after.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop your diarrhea.



- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- If you have diabetes, keep good control of your blood sugar level. Tell your care team if your glucose levels are higher or lower than normal.
- Keeping your pain under control is important to your wellbeing. Please tell your care team if you are experiencing pain.

## Food and Medicine Interactions

- There are no known interactions of lutetium Lu 177 dotatate with food.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with lutetium Lu 177 dotatate. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Wheezing and/or trouble breathing
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Skin flushing (skin feels warm and/or is red)
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your abdomen that does not go away
- Abnormal blood sugar
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Decreased or very dark urine



- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Signs of low potassium levels: weakness, tiredness, muscle cramps, constipation

## Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 7 months after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 4 months after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 2.5 months after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised December 2025

