



Margetuximab-cmkb (Margenza)

About This Drug

Margetuximab-cmkb is used to treat cancer. It is given in the vein (IV).

Possible Side Effects

- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Constipation (unable to move your bowels)
- Pain in your abdomen
- Tiredness and weakness
- Fever
- While you are getting this drug in your vein (IV), you may have a reaction to the drug. Sometimes you may be given medication to stop or lessen these side effects. Your nurse will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Decreased appetite (decreased hunger)
- Effects on the nerves called peripheral neuropathy. You may feel numbness, tingling, or pain in your hands and feet. It may be hard for you to button your clothes, open jars, or walk as usual. The effect on the nerves may get worse with more doses of the drug. These effects get better in some people after the drug is stopped but it does not get better in all people.
- Headache
- Joint and muscle pain
- Pain in your extremities, such as your arms and legs
- Cough
- Trouble breathing
- Hand-foot syndrome. The palms of your hands or soles of your feet may tingle, become numb, painful, swollen, or red.
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in greater than 10% of patients treated with margetuximab-cmkb. All possible side effects are not included. Your side effects may be different



depending on your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Changes in your heart's ability to pump blood properly
- Severe infusion-related reactions

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise. If you feel too tired to exercise vigorously, try taking a short walk.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- If you are not able to move your bowels, check with your doctor or nurse before you use enemas, laxatives, or suppositories.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea and/or constipation.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- Infusion reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.



- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- Avoid sun exposure and apply sunscreen routinely when outdoors.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.

Food and Drug Interactions

- There are no known interactions of margetuximab-cmkb with food.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with margetuximab-cmkb. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and/or weakness that interferes with your daily activities
- Extreme weakness that interferes with normal activities
- Feeling dizzy or lightheaded
- A headache that does not go away
- Pain in your abdomen that does not go away
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Cough that is bothersome
- Wheezing and/or trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- No bowel movement in 3 days or when you feel uncomfortable
- Pain in your abdomen that does not go away
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of the hands, feet, or any other part of the body



- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Pain that does not go away or is not relieved by prescribed medicine
- Painful, red, or swollen areas on your hands or feet
- Numbness, tingling, or pain in your hands and feet
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** This drug may have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 4 months after stopping treatment. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** It is not known if this drug passes into breast milk. For this reason, women should not breastfeed during treatment and for 4 months after stopping treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised March 2023

