

## Melphalan (tablet)

---

### About This Medicine

Melphalan (tablet) is used to treat cancer. It is given orally (by mouth).

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Changes in your liver function
- Thickening and/or inflammation (swelling) of the lung tissues. You may have a cough and/or trouble breathing.
- Rash
- Vasculitis – inflammation of the blood vessels that may cause them to become thick, narrow, scarred, or weak
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

**Note:** All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your medical team.

### Warnings and Precautions

- Severe bone marrow suppression
- Allergic reactions, including anaphylaxis, are rare but may happen in some patients. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**

- This medicine may raise your risk of getting a second cancer, such as myeloproliferative syndrome and acute leukemia.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

## Important Information

- Talk to your doctor before receiving any vaccinations during your treatment. Some vaccinations are not recommended while receiving melphalan (tablet).

## How to Take Your Medicine

- Take this medicine by mouth without food, at least 1 hour before you eat or 2 hours after you eat.
- **Missed dose:** If you vomit or miss a dose, contact your doctor for further instructions. Do not take 2 doses at the same time or extra doses.
- **Handling:** Wash your hands with soap and water before and after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- Cytotoxic medicines leave the body through urine and stool, but they can also be present in other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your health care provider after your treatment. If you take a cytotoxic pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the refrigerator, between 36°F to 46°F (2°C to 8°C). Protect from light.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.



- If you throw up or have diarrhea you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.

## Food and Medicine Interactions

- There are no known interactions of melphalan (tablet) with food, however this medicine should be taken on an empty stomach.
- This medicine may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day



- Pain in your mouth or throat that makes it hard to eat or drink
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Pain in your chest
- Dry cough
- Wheezing and/or trouble breathing
- New rash and/or itching or a rash that is not relieved by prescribed medicines
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- If you think you may be pregnant

## Reproduction Warnings

- **Pregnancy warning:** This medicine may have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment. Let your doctor know right away if you think you may be pregnant.
- In women, menstrual bleeding may become irregular or stop while you are getting this medicine. Do not assume that you cannot become pregnant if you do not have a menstrual period.
- **Breastfeeding warning:** It is not known if this medicine passes into breast milk. For this reason, women should talk to their doctor about the risks and benefits of breastfeeding during treatment with this medicine because this medicine may enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** In men and women both, this medicine may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

New August 2023

