

Mitomycin

About This Medicine

MITOMYCIN (mye toe MYE sin) treats stomach cancer and pancreatic cancer. It works by slowing down the growth of cancer cells. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Fever
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea and vomiting (throwing up)
- Decreased appetite (decreased hunger)
- Skin and tissue irritation including redness, pain, warmth, or swelling at the IV site if the medicine leaks out of the vein and into nearby tissue. Very rarely it may cause local tissue necrosis (death). Skin irritation can happen weeks or months after the infusion.
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe bone marrow suppression, which can be life-threatening.
- Changes in your kidney function
- A syndrome that affects your red blood cells, platelets and blood vessels in your kidneys, which can cause kidney failure and be life-threatening.

- Inflammation (swelling) of the lungs, which may be life-threatening. You may have a dry cough or trouble breathing.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- Drink enough fluids to keep your urine pale yellow. More may be recommended by your care team.
- If you throw up, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.



- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your care team and they can help you get in touch with programs in your local area.

Food and Medicine Interactions

- There are no known interactions of mitomycin with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Wheezing and/or trouble breathing
- Pain in your chest
- Dry cough
- Pain in your mouth or throat that makes it hard to eat or drink
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Swelling of the face, hands, feet, or any other part of the body
- Decreased or very dark urine
- While you are getting this medicine, or afterward, please tell your care team right away if you have any pain, redness, or swelling at the site of the IV infusion.

Reproduction Warnings

- **Pregnancy warning:** Talk with your care team if you are pregnant or planning to become pregnant while taking this medicine. Tell your care team right away if you think you might be pregnant.
- **Breastfeeding warning:** Talk with your care team about breastfeeding during treatment. You may need to stop breastfeeding.



- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Reviewed March 2026

