

## Nivolumab and relatlimab-rmbw (Opdualag)

---

### About This Medicine

NIVOLUMAB; RELATLIMAB (nye VOL ue mab; rel AT li mab) treats skin cancer. It works by helping your immune system slow or stop the spread of cancer cells. It is a monoclonal antibody. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

### Possible Side Effects

- Decrease in white blood cells and red blood cells. This may raise your risk of infection and make you feel more tired and weak.
- Diarrhea (loose bowel movements)
- Tiredness
- Changes in your liver function
- Decreased level of sodium in your blood
- Muscle and bone pain
- Rash and itching

**Note:** Each of the side effects above was reported in 20% or greater of people treated with nivolumab and relatlimab-rmbw. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

### Warnings and Precautions

- This medicine works with your immune system and can cause inflammation (swelling) in any of your organs and tissues and can change how they work. This may put you at risk for developing serious medical problems which can be life-threatening. These side effects may require treatment with steroids at the discretion of your care team.
- Inflammation of the lungs which can be life-threatening. You may have a dry cough or trouble breathing.
- Colitis, which is inflammation in the colon. The symptoms are diarrhea, stomach cramping, and sometimes blood in the bowel movements.
- Changes in your central nervous system can happen. The central nervous system is made up of your brain and spinal cord. You could feel extreme tiredness, agitation, confusion, hallucinations

(see or hear things that are not there), trouble understanding or speaking, loss of control of your bowels or bladder, eyesight changes, numbness or lack of strength to your arms, legs, face, or body, and coma. **If you start to have any of these symptoms let your care team know right away.**

- Severe changes to your liver function, which can cause liver failure and be life-threatening.
- Inflammation in the tissue of the heart
- This medicine may affect some of your hormone glands (especially the thyroid, adrenals, pituitary, and pancreas).
- Blood sugar levels may change, and you may develop diabetes. If you already have diabetes, changes may need to be made to your diabetes medicine.
- Changes in your kidney function
- Allergic skin reaction, which can very rarely be life-threatening. You may develop blisters on your skin that are filled with fluid or a severe red rash all over your body that may be painful. Sometimes, a serious and life-threatening reaction to this medicine can happen that can cause a skin rash, along with problems with your organs, swollen lymph nodes, fever and higher than normal white blood cells.
- While you are getting this medicine in your vein (IV), you may have a reaction to the medicine. Sometimes you may be given medicine to stop or lessen these side effects. Your care team will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Increased risk of serious complications that can be life-threatening, such as graft versus host disease (GVHD), in patients who undergo a stem cell transplant before or after receiving nivolumab and relatlimab-rmbw.
- Increased risk of organ rejection in people who have received donor organs.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

## Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.



- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- Drink enough fluids to keep your urine pale yellow.
- If you have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid). Eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that is available to help stop or lessen diarrhea.
- If you have diabetes, keep good control of your blood sugar level. Tell your care team if your glucose levels are higher or lower than normal.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- To help with itching, moisturize your skin several times a day.
- Avoid sun exposure and apply sunscreen routinely when outdoors.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.
- Infusion reactions may happen after your infusion. **If this happens, call 911 for emergency care.**

## Food and Medicine Interactions

- There are no known interactions of nivolumab and relatlimab-rmbw with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call Your Care Team

Not all possible side effects are included. Some of these side effects, although rare, can be life-threatening.

### Lung problems:

- Inflammation of the lungs
- Cough
- Trouble breathing
- Upper respiratory tract infection

### Call your care team if you have any of these symptoms:

- Wheezing or trouble breathing



- New or worsening cough
- Chest pain
- Coughing up yellow, green, or bloody mucus

### **Heart Problems**

- Inflammation in the tissue of the heart

### **Call your care team if you have any of these symptoms:**

- Trouble breathing
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Pain in your chest
- Swelling of the hands, feet, or any other part of the body

### **Stomach problems:**

- Nausea
- Decreased appetite (decreased hunger)
- Diarrhea
- Constipation (unable to move bowels)
- Pain in your abdomen
- Inflammation of your colon
- Blood in your stool

### **Call your care team if you have any of these symptoms:**

- Nausea that stops you from eating or drinking or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- No bowel movement for 3 days or when you feel uncomfortable
- Pain in your abdomen that does not go away
- Blood in your stool (bright red or black/tarry)

### **Liver problems:**



- Changes in your liver function

**Call your care team if you have any of these symptoms:**

- Yellowing of the eyes or skin
- Dark urine
- Pale bowel movements
- Pain on the right side of your abdomen that does not go away
- Feeling very tired and weak
- Unusual itching
- Easy bleeding or bruising

**Hormone gland problems:**

- Changes in some of your hormone glands (especially the thyroid, adrenals, pituitary, and pancreas)
- Blood sugar levels may change, and you may develop diabetes

**Call your care team if you have any of these symptoms:**

- Headache that does not go away
- Tiredness and weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Changes in mood or behavior such as irritability and/or feeling forgetful
- Shakiness
- Weight loss or weight gain
- Nausea
- Abnormal blood sugar
- Unusual thirst or passing urine often
- Feeling cold

**Kidney problems:**

- Changes in your kidney function

**Call your care team if you have any of these symptoms:**

- Decreased or very dark urine
- Cloudy urine and/or urine that smells bad
- Trouble urinating
- Pain or burning when you pass urine
- Feeling like you have to pass urine often, but not much comes out when you do
- Tender or heavy feeling in your lower abdomen
- Pain on one side of your back under your ribs

**Skin problems:**

- Rash and itching
- Allergic skin reaction
- Soreness of the mouth and throat

**Call your care team if you have any of these symptoms:**

- New rash and/or itching
- Fluid-filled bumps/blisters
- Rash that is not relieved by prescribed medicines
- Red areas, white patches, or sores in your mouth that hurt
- Flu-like symptoms: fever, headache, muscle and joint aches, and tiredness
- Swollen lymph nodes in your neck and/or armpits

**Inflammation of the brain:**

- Changes in your brain and spinal cord
- Headache
- Effects on the nerves

**Call your care team if you have any of these symptoms:**

- Headache that does not go away
- Extreme tiredness, agitation, or confusion
- Seizures
- Hallucinations
- Trouble understanding or speaking
- Loss of control of bowels or bladder

- Numbness or lack of strength to your arms, legs, face, or body
- Numbness, tingling, pins, and needles, or pain in your arms, hands, legs, or feet

### Other problems:

- Low red blood cells, and platelets
- Fever
- Inflammation of your eye and/or other changes in vision
- Allergic reaction to the medicine
- Electrolyte changes
- Muscle, bone, and joint pain

### Call your care team if you have any of these symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills, flushing
- Easy bleeding or bruising
- Blurred vision or other changes in eyesight
- Sensitivity to light
- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**
- Pain that does not go away, or is not relieved by prescribed medicines
- Extreme muscle weakness

### Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for at least 5 months after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for at least 5 months after the last dose.
- **Fertility warning:** The effect of this medicine on fertility is not known. If you plan to have children, talk with you care team.

Revised July 2025

