

Panobinostat (Farydak)

About This Drug

Panobinostat is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Fever
- Tiredness
- Swelling of your legs, ankles and/or feet
- Decreased appetite (decreased hunger)
- Electrolyte changes
- Decrease in a blood protein called albumin
- Changes in your kidney and liver function

Note: Each of the side effects above was reported in 20% or greater of patients treated with panobinostat. Not all possible side effects are included above.

Warnings and Precautions

- Severe bone marrow suppression
- Severe and life-threatening infections
- Abnormal electrocardiogram (EKG/ECG) and risk of heart attack
- Abnormal bleeding, which can be life-threatening – symptoms may be coughing up blood, throwing up blood (may look like coffee grounds), red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding.
- Severe diarrhea

Important Information



- Your doctor will recommend medication to decrease your diarrhea. Please call your doctor or nurse if you have diarrhea.

How to Take Your Medication

- Swallow the medicine whole, with a cup of water, with or without food at the same time each day it is scheduled. Do not chew, crush, or open the capsules.
- **Missed dose:** If you miss a dose, take it as soon as you think about it. If it is within 12 hours of your next dose, then skip the missed dose and contact your doctor. Do not take 2 doses at the same time and do not double up on the next dose.
- If you vomit a dose, take your next dose at the regular time. Do not take 2 doses at the same time.
- **Handling:** Wash your hands after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- If any of the capsules are broken, do not touch them with bare hands. Carefully throw away the capsules and wash your hands after handling.
- If you get any of the content of a broken capsules on your skin or in your eyes, you should wash the area of the skin well with soap and water right away. Call your doctor if you get a skin reaction.
- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- **Storage:** Store this medicine in the original container at room temperature. Protect from light.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.



- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have diarrhea you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.

Food and Drug Interactions

- Avoid grapefruit, star fruit, pomegranate, grapefruit juice and/or pomegranate juice while taking this medicine as it may raise the levels of panobinostat in your body which could make side effect worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with panobinostat. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- Avoid the use of St. John's Wort while taking panobinostat as this may lower the levels of the drug in your body, which can make it less effective.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness or weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Feeling that your heart is beating in a fast or not normal way (palpitations)
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back or it can be constant. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Swelling of your legs, ankles and/or feet
- Weight gain of 5 pounds in one week (fluid retention)
- Decreased or very dark urine



- Signs of abnormal bleeding such as coughing up blood, throwing up blood (may look like coffee grounds), easy bleeding or bruising, red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 3 months after treatment. Men with female partners of childbearing potential should use condoms during your cancer treatment and for 6 months after your cancer treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** It is not known if this drug passes into breast milk. For this reason, women should not breastfeed during treatment with this drug because this drug may enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** In men and women both, this drug may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised June 2021

