

## Pirtobrutinib (Jaypirca)

---

### About This Medicine

PIRTOBRUTINIB (PIR toe BROO ti nib) treats leukemia and lymphoma. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions.

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Diarrhea (loose bowel movements)
- Tiredness
- Muscle and bone pain
- Bruising
- Cough
- COVID-19 infection
- Changes in your kidney function

**Note:** Each of the side effects above was reported in 20% or greater of people treated with pirtobrutinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

### Warnings and Precautions

- Risk of severe and life-threatening infections
- Abnormal bleeding, which can be life-threatening – symptoms may be coughing up blood, throwing up blood (may look like coffee grounds), red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding.
- Abnormal heartbeat
- Severe bone marrow suppression

- This medicine may raise your risk of getting a second cancer, such as skin cancer or other solid tumors.
- Changes in your liver function, which can cause liver failure and be life-threatening.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

## Important Information

- You may need to hold pirtobrutinib for 3 to 7 days before and after any planned medical, surgical and/or dental procedures. Talk to your care team about any precautions you may need to take.

## How to Take Your Medication

- Swallow the medicine whole with water, with or without food. Do not chew, cut, or crush tablets.
- Take it at approximately the same time each day.
- **Missed dose:** If you miss a dose by less than 12 hours, take the missed dose right away and go back to your normal schedule. If you miss a dose by more than 12 hours, then skip the missed dose and go back to your normal schedule.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.



- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid). Eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop diarrhea.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- Avoid sun exposure and apply sunscreen routinely when outdoors. Cover up when you are out in the sun. Wear wide-brimmed hats, long-sleeved shirts, and pants.

## Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your care team as this could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with pirtobrutinib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

## When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness or extreme weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Wheezing and/or trouble breathing
- A cough that is bothersome
- New loss of taste or smell
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Decreased urine, or very dark urine
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin

- Signs of abnormal bleeding such as coughing up blood, throwing up blood (may look like coffee grounds), easy bleeding or bruising, red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding
- Pain that does not go away or is not relieved by prescribed medicine
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, difficulty urinating, burning or pain when you pass urine, redness and/or swelling of the skin

## Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 1 week after the last dose. Tell your care team right away if you think you might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 week after the last dose.
- **Fertility:** The effect of this medicine on fertility is not known. If you plan to have children, talk with your care team.

Revised August 2025

