

## Pirtobrutinib (Jaypirca)

---

### About This Medicine

Pirtobrutinib is used to treat cancer. It is given orally (by mouth).

### Possible Side Effects

- Diarrhea (loose bowel movements)
- Tiredness
- Muscle and bone pain
- Bruising
- Cough
- COVID-19 infection

**Note:** Each of the side effects above was reported in 20% or greater of patients treated with pirtobrutinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your medical team.

### Warnings and Precautions

- Risk of severe and life-threatening infections
- Abnormal bleeding, which can be life-threatening – symptoms may be coughing up blood, throwing up blood (may look like coffee grounds), red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding.
- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Abnormal heartbeat
- This medicine may raise your risk of getting a second cancer, such as skin cancer or other solid tumors.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

## Important Information

- You may need to hold pirtobrutinib for 3 to 7 days before and after any planned medical, surgical and/or dental procedures. Talk to your doctor and/or nurse about any precautions you may need to take.

## How to Take Your Medication

- Swallow the medicine whole with water, with or without food. Do not chew, cut, or crush it.
- Take at approximately the same time each day.
- **Missed dose:** If you miss a dose by less than 12 hours, take the missed dose right away and go back to your normal schedule. If you miss a dose by more than 12 hours, then skip the missed dose and go back to your normal schedule.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- Cytotoxic medicines leave the body through urine and stool, but they can also be present in other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your health care provider after your treatment. If you take a cytotoxic pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your doctor's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid). Eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.



- Ask your nurse or doctor about medicine that can lessen or stop diarrhea.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.
- Avoid sun exposure and apply sunscreen routinely when outdoors.

## Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your doctor as this could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with pirtobrutinib. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

## When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness or extreme weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Wheezing and/or trouble breathing
- A cough that is bothersome
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Signs of abnormal bleeding such as coughing up blood, throwing up blood (may look like coffee grounds), easy bleeding or bruising, red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding
- Pain that does not go away or is not relieved by prescribed medicine
- Signs of infection: fever or chills, cough, trouble breathing, severe pain in your abdomen, difficulty urinating, burning or pain when you pass urine, redness and/or swelling of the skin
- If you think you may be pregnant

## Reproduction Warnings

- **Pregnancy warning:** This medicine can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment



and for 1 week after stopping treatment. Let your doctor know right away if you think you may be pregnant.

- **Breastfeeding warning:** Women should not breast feed during treatment and for 1 week after stopping treatment because this medicine could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this medicine. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised January 2024

