

Rho(D) Immune Globulin (WinRho SDF, Rhophylac)

About This Medicine

RhO [D] IMMUNE GLOBULIN (ROE D im MYOON GLOB you lin) treats low levels of platelets in your body caused by immune thrombocytopenia (ITP). It may also be used to help prevent your immune system from reacting if you have Rh-negative blood and come into contact with Rh-positive blood. This can happen during a blood transfusion. It can also happen during pregnancy if the fetus has a different blood type than you. It works by helping your body ignore the Rh protein. It is given in the vein (IV) or as an injection in a muscle (IM).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Fever and chills
- Weakness
- Headache
- Feeling dizzy
- Breakdown of your red blood cells, which can cause anemia (decreased red blood cells). This breakdown of red blood cells can be life-threatening.

Note: Each of the side effects above was reported in 5% or greater of people treated with Rho(D) immune globulin. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Allergic reactions, including anaphylaxis, are rare but may happen in some people. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**
- Fluid may build-up around your lungs following treatment. You may have a fever, dry cough or trouble breathing.
- You may be at risk of blood clots and events such as stroke and heart attack. A blood clot in your leg may cause your leg to swell, appear red and warm, and/or cause pain. A blood clot in your lungs may cause trouble breathing, pain when breathing, and/or chest pain.

- Changes in your kidney function, which can cause kidney failure and may be life-threatening.
- Rho(D) immune globulin is made from human plasma and carries a very rare risk of transmitting infectious diseases.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- Talk to your care team before receiving any vaccinations during your treatment. Some vaccinations are not recommended while receiving Rho(D) immune globulin or within 3 months of treatment.
- Rho(D) immune globulin may interfere with certain blood tests and some type of blood sugar monitoring systems, which can falsely increase your blood sugar readings. Please talk to your care team for more information.

Treating Side Effects

- If you are dizzy, get up slowly after sitting or lying.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.

Food and Medicine Interactions

- There are no known interactions of Rho(D) immune globulin with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Headache that does not go away
- Feeling dizzy or lightheaded
- Wheezing and/or trouble breathing
- Dry cough
- Extreme weakness that interferes with normal activities



- Your leg is swollen, red, warm and/or painful
- Weight gain of 5 pounds in one week (fluid retention)
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back, or it can be constant. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Symptoms of a stroke such as sudden numbness or weakness of your face, arm, or leg, mostly on one side of your body; sudden confusion, trouble speaking or understanding; sudden trouble seeing in one or both eyes; sudden trouble walking, feeling dizzy, loss of balance or coordination; or sudden, bad headache with no known cause. **If you have any of these symptoms for 2 minutes, call 911.**
- Easy bleeding and/or bruising
- Decreased urine, blood in your urine, or very dark urine
- Severe back pain
- Abnormally high blood sugar levels on home blood sugar monitoring systems
- Yellowing of the eyes or skin
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**

Reproduction Warnings

- **Pregnancy warning:** Talk with your care team if you are pregnant or planning to become pregnant while taking this medicine. Tell your care team right away if you think you might be pregnant.
- **Breastfeeding warning:** Talk with your care team about breastfeeding during treatment. You may need to stop breastfeeding.

Revised March 2026

https://winrho.com/pdfs/kamada_winrho-uspi.pdf

<https://labeling.cslbehrling.com/PI/US/Rhophylac/EN/Rhophylac-Prescribing-Information.pdf>

Disclaimer

User/Authorized User acknowledges that the ClinicalPath Portal is intended to be utilized as an information management tool only, and that Elsevier has not represented the ClinicalPath Portal as having the ability to diagnose disease, prescribe treatment, or perform any other tasks that constitute the practice of providing medical care. The clinical information contained within the ClinicalPath Portal is intended as a supplement to, and not a substitute for, the knowledge, expertise, skill, and judgment of physicians, pharmacists and other healthcare professionals involved with patient care at User/Authorized User facilities. CONFIDENTIAL AND PROPRIETARY. NOT FOR DISTRIBUTION.