

Ripretinib (Qinlock)

About This Medicine

RIPRETINIB (rip RE ti nib) treats gastrointestinal stromal tumor (GIST). It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Constipation (not able to move bowels)
- Pain in your abdomen
- Tiredness
- Decreased appetite (decreased hunger)
- Muscle pain/aching
- Hand-foot syndrome. The palms of your hands or soles of your feet may tingle, become numb, painful, swollen, or red.
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in 20% or greater of people treated with ripretinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- This medicine may raise your risk of getting new skin cancers.
- High blood pressure, which can be severe.
- Congestive heart failure – your heart is not pumping blood as well as it should be, and fluid can build up in your body.
- Slow wound healing



- Sensitivity to sunlight/light. You may get a skin rash/reaction if you are in the sun or are exposed to sun lamps and tanning beds.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- Ripretinib may cause slow wound healing. It should not be given within at least 1 week of planned surgery and for at least 2 weeks after major surgical procedures and until the wound is fully healed. If you must have emergency surgery or have an accident that results in a wound, tell the care team that you are on ripretinib.

How to Take Your Medicine

- Swallow the medicine whole with or without food. Take this medicine at the same time each day.
- **Missed dose:** If you miss a dose by more than 8 hours, then skip the missed dose and go back to your normal schedule. If it has been less than 8 hours since your missed dose, then take the missed dose. Do not take 2 doses at the same time or extra doses. Do not replace a vomited dose.
- If you are taking a class of medicine called CYP3A inducer (e.g., efavirenz) while taking ripretinib, it may affect how you handle a missed dose of ripretinib. Check with your care team about any special instructions you may need to follow.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature with the desiccant to protect from moisture and light. Replace cap securely each time after opening. Do not remove desiccant. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- Drink enough fluids to keep your urine pale yellow.



- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can help stop or lessen your diarrhea, constipation and/or nausea/vomiting.
- If you are not able to move your bowels, check with your care team before you use enemas, laxatives, or suppositories.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- Avoid sun exposure during treatment and for at least 1 week after stopping treatment with ripretinib. Use sunscreen with SPF 30 or higher when you are outdoors even for a short time. Cover up when you are out in the sun. Wear wide-brimmed hats, long-sleeved shirts, and pants. Keep your neck, chest, and back covered.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- Use a gentle, unscented lotion to keep your hands and feet soft. A cream with 10% or 20% urea may help. Don't rub your hands or feet too much.
- Try not to put your hands or feet in hot water, like baths, showers, or washing dishes. Wear gloves when doing work with your hands. Avoid tight shoes or socks.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your care team and they can help you get in touch with programs in your local area.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.

Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your care team as this could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with ripretinib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.



- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- A headache that does not go away
- Blurred vision or other changes in eyesight
- Trouble breathing
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Tiredness and/or extreme weakness that interferes with your daily activities
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- No bowel movement in 3 days or when you feel uncomfortable
- Pain in your abdomen that does not go away
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Swelling of the legs, feet, or ankles
- Weight gain of 5 pounds in one week (fluid retention)
- Painful, red, or swollen areas on your hands or feet
- Numbness and/or tingling of your hands and/or feet
- Develop sensitivity to sunlight/light
- A new rash or a rash that is not relieved by prescribed medicines
- New skin sore, or reddish bump that bleeds or does not heal
- New skin lesion and/or wart
- Change in size or color of a mole

Reproduction Warnings

- **Pregnancy warning:** This medicine may cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 1 week after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 1 week after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 week after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.



Revised January 2026

