

Ropeginterferon alfa-2b-njft (Besremi)

About This Medicine

ROPEGINTERFERON ALFA-2B (ROE peg IN ter FEER on al fa 2b) treats blood and bone marrow cancers. It works by slowing down the growth of cancer cells. It is given as an injection under the skin (subcutaneously).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Flu-like symptoms: fever, chills, headache, muscle/joint aches, and tiredness
- Bone, muscle and/or joint pain
- Tiredness
- Itching
- Inflammation of the nasal passages and throat

Note: Each of the side effects above was reported in greater than 40% of people treated with ropeginterferon alfa-2b-njft. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe depression and other psychiatric disorders such as aggression, feeling irritable and/or abnormally well, confusion, thoughts of hurting yourself, and suicide.
- Changes in your thyroid function and blood sugar levels
- Heart changes such as abnormal heartbeat, changes in your heart's ability to pump blood properly which can lead to fluid build-up in your body, and/or risk of heart attack.
- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Allergic reactions, including anaphylaxis, are rare but may happen in some people. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**
- Inflammation (swelling) of your pancreas

- Severe colitis, which is inflammation in the colon and can be life-threatening. Symptoms are diarrhea (loose bowel movements), stomach cramping, and sometimes blood in the bowel movements.
- Changes in your lungs, including inflammation and/or scarring of the lungs, increased blood pressure in the arteries of your lungs, and infection. You may have a cough and/or trouble breathing. Some changes in your lungs can be life-threatening.
- Blurred vision or other changes in eyesight such as problems with your retina (thin membrane that covers the back of the eye) which can very rarely cause blindness (loss of vision).
- Increase in your cholesterol and/or triglyceride levels
- Changes in your liver function
- Changes in your kidney function
- Tooth and gum problems which may cause tooth loss. You may have dry mouth that can damage your teeth.
- Skin problems such as rash, itching, thickening of the skin and hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- This medicine may impair your ability to drive or use machinery. Use caution and talk your care team about any precautions you may need to take.

How to Take Your Medicine

- Talk to your care team for proper preparation, dosing, and administration if you are self-injecting this medicine.
- **Missed dose:** If you miss a dose, contact your care team right away.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the refrigerator, between 36°F to 46°F (2°C to 8°C). Store in original carton to protect from light. Keep away from heat. Do not freeze. Discuss with your care team how to dispose of unused medicine/needles. Keep out of the reach of children and pets.



Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your care team before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid). Eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that is available to help stop or lessen diarrhea.
- If you have diabetes, keep good control of your blood sugar level. Tell your care team if your glucose levels are higher or lower than normal.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your care team and they can help you get in touch with programs in your local area.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- To help with itching, moisturize your skin several times a day.
- Avoid sun exposure and apply sunscreen routinely when outdoors.
- Sugar-free hard candies and chewing gum can keep your mouth moist.
- Tell your dentist that you are taking this medicine. Talk to your care team about any dental problems you may have before starting this medicine. You may need to see your dentist to have a dental exam and fix any dental problems before starting this medicine. Take good care of your teeth and brush two times a day. Make sure you see your dentist for regular follow-up appointments.

Food and Medicine Interactions

- There are no known interactions of ropeginterferon alfa-2b-njft with food.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this



medicine as there are known medicine interactions with ropeginterferon alfa-2b-njft. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness or extreme weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Blurred vision or other changes in eyesight
- Flashes of light and/or dark areas in your field of vision, colored dots, or halos
- Loss of vision
- Confusion
- Severe mood changes such as depression or unusual thoughts and/or behaviors
- Lose interest in your daily activities that you used to enjoy and feel this way every day, and/or you feel hopelessness
- Thoughts of hurting yourself or others, and suicide
- Feeling irritable, nervous, or restless
- Feeling abnormally well
- Easy bleeding or bruising
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back, or it can be constant. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Wheezing and/or trouble breathing
- Lips or skin turn a bluish color
- Dry cough
- Coughing up yellow, green, or bloody mucus
- Tooth and/or gum pain
- Difficulty eating/swallowing
- Sore mouth/throat
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Severe pain in your abdomen that may spread to your back
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Blood in your stool (bright red, or black/tarry)
- Weight gain of 5 pounds in one week (fluid retention)
- Abnormal blood sugar



- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Pain that does not go away, or is not relieved by prescribed medicines
- Swelling of the hands, feet, or any other part of the body
- Decreased or very dark urine
- A new rash or a rash that is not relieved by prescribed medicines
- Itching that is bothersome
- Signs of low thyroid function: tiredness, unexplained weight gain, hair loss, dry skin, constipation, increased sensitivity to cold
- Signs of increased thyroid function: unexplained weight loss, sweating, irritability, feeling that your heart is beating fast or in a not normal way (palpitations)
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Flu-like symptoms: fever, headache, muscle/joint aches, and tiredness

Reproduction Warnings

- **Pregnancy warning:** This medicine may cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for at least 8 weeks after the last dose. Tell your care team right away if you think you might be pregnant.
- Menstrual bleeding may become irregular or stop while you are getting this medicine. Do not assume that you cannot become pregnant if you do not have a menstrual period.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 8 weeks after the last dose.
- **Fertility warning:** The effect of this medicine on fertility is not known. If you plan to have children, talk with you care team.

Revised March 2026

Disclaimer

User/Authorized User acknowledges that the ClinicalPath Portal is intended to be utilized as an information management tool only, and that Elsevier has not represented the ClinicalPath Portal as having the ability to diagnose disease, prescribe treatment, or perform any other tasks that constitute the practice of providing medical care. The clinical information contained within the ClinicalPath Portal is intended as a supplement to, and not a substitute for, the knowledge, expertise, skill, and judgment of physicians, pharmacists and other healthcare professionals involved with patient care at User/Authorized User facilities. CONFIDENTIAL AND PROPRIETARY. NOT FOR DISTRIBUTION.