

Sacituzumab govitecan-hziy (Trodelvy)

About This Medicine

Sacituzumab govitecan-hziy is used to treat cancer. It is given in your veins (IV).

Possible Side Effects

- Decrease in the number of white blood cells and red blood cells. This may raise your risk of infection and make you feel more tired.
- Nausea and vomiting (throwing up)
- Constipation (unable to move your bowels)
- Diarrhea (loose bowel movements)
- Tiredness
- Decreased appetite (decreased hunger)
- Blood sugar levels may change
- Decrease in a blood protein called albumin
- Electrolyte changes
- Changes in your kidney function
- Changes in your liver function
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in 25% or greater of patients treated with sacituzumab govitecan-hziy. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Severe decrease in white blood cells and neutropenic fever, which is a type of fever that can develop when you have a very low number of white blood cells and can be life-threatening
- Severe nausea and vomiting
- Severe diarrhea and/or inflammation (swelling) in the colon in the setting of severely low white blood cells, which raises your risk of infection.

- Allergic reactions, including anaphylaxis are rare but may happen in some patients, and may be life-threatening. Signs of allergic reaction to this medicine may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this medicine. You should get urgent medical treatment.**
- While you are getting this medicine in your vein (IV), you may have a reaction to the medicine. Sometimes you may be given medicine to stop or lessen these side effects. Your nurse will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- People that carry a gene for uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1)*28 allele have an increased risk for side effects, especially low white blood cells, neutropenic fever, and low red blood cells. Talk to your doctor if you have questions, or have ever been told that you carry this gene.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Important Information

- Cytotoxic medicines leave the body through urine and stool, but they can also be present in other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your health care provider after your treatment. If you take a cytotoxic pill each day, follow these precautions every day.
- It is important that you notify your doctor and/or nurse at the first sign of diarrhea so they can provide you with anti-diarrhea medicine and give you further instructions. Notify your doctor and/or nurse if you are taking anti-diarrhea medicine and your symptoms have not improved or are worsening after 24 hours.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- Drink enough fluids to keep your urine pale yellow.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.



- If you are not able to move your bowels, check with your doctor or nurse before you use enemas, laxatives, or suppositories.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea, constipation, and/or nausea.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- If you have diabetes, keep good control of your blood sugar level. Tell your nurse or your doctor if your glucose levels are higher or lower than normal.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.
- Infusion reactions may occur after your infusion. **If this happens, call 911 for emergency care.**

Food and Medicine Interactions

- There are no known interactions of sacituzumab govitecan-hziy with food.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with sacituzumab govitecan-hziy. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and/or weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Blood in your stool, or black stools
- No bowel movement in 3 days or when you feel uncomfortable



- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Decreased or very dark urine
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Swelling of the hands, feet, or any other part of the body
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This medicine can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 6 months after stopping treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for 3 months after stopping treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 1 month after stopping treatment because this medicine could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** In women, this medicine may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on egg banking.

Revised April 2024

