

Sonidegib (Odomzo)

About This Medicine

SONIDEGIB (soe ni DEG ib) treats skin cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Pain in your abdomen
- Tiredness
- General pain
- Weight loss
- Decreased appetite (decreased hunger)
- Bone and muscle pain/aching
- Muscle spasm
- Changes in the way food and drinks taste
- Headache
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.
- Itching

Note: *The side effects above were reported in 10% or greater of people treated with sonidegib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.*

Warnings and Precautions



- Muscle problems and rhabdomyolysis - damage to your muscles which may release proteins in your blood and affect how your kidneys work, which may be life-threatening. You may have severe muscle weakness and/or pain, or dark urine.

Note: *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.*

Important Information

- Do not donate blood during your treatment and for at least 20 months after treatment.
- Do not donate sperm during your treatment and for at least 8 months after treatment because this medicine is present in semen.

How to Take Your Medicine

- Swallow the medicine whole on an empty stomach, at least 1 hour before you eat or 2 hours after you eat.
- **Missed dose:** If you miss a dose, take your next dose at the regular time. Do not take 2 doses at the same time and do not double up on the next dose.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your care team's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).



- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your care team about medicine that can lessen or stop your diarrhea.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your care team's approval, may increase your appetite.
- To help with weight loss, drink fluids that contribute calories (whole milk, juice, soft drinks, sweetened beverages, milkshakes, and nutritional supplements) instead of water.
- Keeping your pain under control is important to your well-being. Please tell your care team if you are experiencing pain.
- Taking good care of your mouth may help food taste better and improve your appetite.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your care team and they can help you get in touch with programs in your local area.
- To help with itching, moisturize your skin several times a day.
- Avoid sun exposure and apply sunscreen routinely when outdoors.

Food and Medicine Interactions

- This medicine may interact with grapefruit and grapefruit juice. Talk to your care team, as this could make side effects worse.
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with sonidegib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Headache that does not go away
- Tiredness that interferes with your daily activities



- Pain in your abdomen or general pain that does not go away or is not relieved by prescribed medicine
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Changes in the way food and drink taste that is causing a loss of appetite
- Unexplained muscle spasms, pain, tenderness or weakness
- Signs of rhabdomyolysis such as decreased or very dark urine, muscle pain in the shoulders, thighs, or lower back; muscle weakness or trouble moving arms and legs.
- Itching that is bothersome

Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 20 months after the last dose. If you can get your partner pregnant, use a condom, even after a vasectomy while taking this medicine and for 8 months after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 20 months after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

Revised January 2026

