

Tagraxofusp-erzs (Elzonris)

About This Drug

Tagraxofusp-erzs is used to treat cancer. It is given in the vein (IV).

Possible Side Effects

- Decrease in red blood cells and in the number of platelets. This may make you feel more tired and raise your risk of bleeding.
- Nausea
- Fever
- Tiredness
- Swelling of your legs, ankles and/or feet
- Weight gain
- Electrolyte changes
- Decrease in a blood protein called albumin
- Changes in your liver function
- Blood sugar levels may change
- Capillary leak syndrome – a syndrome where fluid and protein can leak from your blood vessels into your tissues. This can cause a decrease in your blood protein level and blood pressure and fluid can accumulate in your tissues and/or lungs.

Note: Each of the side effects above was reported in 30% or greater of patients treated with tagraxofusp-erzs. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Severe and life-threatening capillary leak syndrome
- Allergic reactions, including anaphylaxis are rare but may happen in some patients. Signs of allergic reaction to this drug may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this drug. You should get urgent medical treatment.**

- Severe changes in your liver function

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To help decrease your risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- To help with nausea, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you have diabetes, keep good control of your blood sugar level. Tell your nurse or your doctor if your glucose levels are higher or lower than normal.

Food and Drug Interactions

- There are no known interactions of tagraxofusp-erzs with food.
- This drug may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and/or weakness that interferes with your daily activities



- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Wheezing and/or trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of your legs, ankles and/or feet
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** This drug may have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 1 week after stopping treatment. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 1 week after stopping treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised January 2023

