

Talazoparib (Talzenna)

About This Drug

Talazoparib is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Nausea and vomiting
- Diarrhea (loose bowel movements)
- Tiredness
- Changes in your liver function
- Decreased level of calcium in your blood
- Blood sugar levels may change
- Decreased appetite (decreased hunger)
- Headache
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in 20% or greater of patients treated with talazoparib. Not all possible side effects are included above.

Warnings and Precautions

- This drug may raise your risk of getting a second cancer, such as acute myeloid leukemia and/or myelodysplastic syndrome.
- Severe bone marrow suppression

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

How to Take Your Medication

- Swallow the medicine whole with or without food. Do not open or dissolve capsules.
- **Missed dose:** If you vomit or miss a dose, take your next dose at the regular time, and contact your physician. Do not take 2 doses at the same time and do not double up on the next dose.
- **Handling:** Wash your hands after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- **Storage:** Store this medicine in the original container at room temperature.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- If you have diabetes, keep good control of your blood sugar level. Tell your nurse or your doctor if your glucose levels are higher or lower than normal.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.



- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.

Food and Drug Interactions

- There are no known interactions of talazoparib with food.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with talazoparib. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Headache that does not go away
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Signs of low calcium such as numbness or tingling around your mouth or in your hands and/or feet, muscle stiffness, twitching, spasms, or cramps
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 7 months after treatment. Men with female partners of childbearing potential should



use effective methods of birth control during your cancer treatment and for at least 4 months after your cancer treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.

- **Breastfeeding warning:** Women should not breastfeed during treatment and for at least 1 month after treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** In men, this drug may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm banking.

Revised November 2021

