

## Taletrectinib (Ibtrozi)

---

### About This Medicine

TALETRECTINIB (TA le TREK tin ib) treats lung cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team if you have questions

### Possible Side Effects

- Decreased white blood cells and red blood cells. This may raise your risk of infection and make you tired and weak.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Constipation (not able to move bowels)
- Tiredness
- Feeling dizzy or lightheaded
- Rash
- Electrolyte changes
- Increase in your cholesterol and triglyceride levels
- Increased uric acid in your blood
- Decreased albumin in your blood
- Changes in your kidney function
- Changes in your liver function
- Increase in creatine phosphokinase (CPK), which can sometimes be the result of muscle inflammation or damage. You may have muscle aching and/or cramping.

**Note:** Each of the side effects above was reported in 20% or greater of people treated with taletrectinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

### Warnings and Precautions

- Severe changes in your liver function, which can cause liver failure and be life-threatening.
- Inflammation (swelling) and/or scarring of the lungs, which may be life-threatening. You may have a cough and/or trouble breathing.
- Abnormal electrocardiogram (EKG/ECG), which can be life-threatening
- Increased uric acid in the blood
- Muscle pain or weakness
- Risk of bone fractures

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

## How to Take Your Medicine

- Take this medicine on an empty stomach, at approximately the same time each day. Do not eat food for at least 2 hours before and for 2 hours after taking taletrectinib. Swallow capsules whole. Do not open, chew, crush or dissolve capsules.
- **Missed dose:** If you vomit or miss a dose, take your next dose at the regular time. Do not take 2 doses at the same time and do not double up on the next dose.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your care team about other helpful tips and medicine that is available to help stop or lessen these symptoms.



- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- If you are not able to move your bowels, check with your care team before you use enemas, laxatives, or suppositories.
- Ask your care team about medicine that can lessen or stop diarrhea and/or constipation.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- During treatment and for at least 5 days after your treatment, minimize sun exposure and use sunscreen with SPF 30 or higher when you are outdoors even for a short time. Cover up when you are out in the sun. Wear wide-brimmed hats, long-sleeved shirts, and pants. Keep your neck, chest, and back covered.

## Food and Medicine Interactions

- Avoid grapefruit or grapefruit juice while taking this medicine as it may raise the levels of taletrectinib in your body which could make side effects worse. Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with taletrectinib. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- This medicine may interact with St. John's Wort and may lower the levels of the medicine in your body, which can make it less effective.
- Medicines that treat heartburn and stomach upset may lower the effect of your cancer treatment if taken with taletrectinib. Ask your care team what medicine you can take with taletrectinib to help with heartburn or stomach upset.

## When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and/or weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Pain in your chest
- Dry cough
- Wheezing and/or trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your abdomen that does not go away
- No bowel movement in 3 days or when you feel uncomfortable
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy



- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Swelling of the hands, feet, or any other part of the body
- Unexplained muscle pain, tenderness, or weakness
- Warm, red, tender, or swollen joints, especially your big toe
- Urine that is pink or brownish in color
- Decreased or very dark urine
- Pain that does not go away, or is not relieved by prescribed medicines
- New rash and/or itching or a rash that is not relieved by prescribed medicines

## Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 3 weeks after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 3 weeks after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 3 weeks after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

New July 2025

