

Tazemetostat (Tazverik)

About This Drug

Tazemetostat is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Nausea and vomiting (throwing up)
- Constipation (not able to move bowels)
- Pain in your abdomen
- Tiredness
- Decreased appetite (decreased hunger)
- General pain and pain in your bones and/or muscles
- Upper respiratory tract infection

Note: Each of the side effects above was reported in 20% or greater of patients treated with tazemetostat. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- This drug may raise your risk of getting a second cancer, such as myelodysplastic syndrome and/or acute myeloid leukemia.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

How to Take Your Medication

- Swallow the medicine whole with or without food. Do not cut, crush, or chew tablets.
- **Missed dose:** If you vomit or miss a dose, take your next dose at the regular time. Do not take 2 doses at the same time and do not double up on the next dose.
- **Handling:** Wash your hands after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- **Storage:** Store this medicine in the original container at room temperature. Do not store above 30°C (86°F).
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Ask your doctor or nurse about medicines that are available to help stop or lessen constipation.
- If you are not able to move your bowels, check with your doctor or nurse before you use enemas, laxatives, or suppositories.
- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your doctor's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.

Food and Drug Interactions

- Avoid grapefruit or grapefruit juice while taking this medicine as it may raise the levels of tazemetostat in your body which could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with tazemetostat. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplement to make sure that there are no interactions.
- Avoid the use of St. John's Wort while taking tazemetostat as this may lower the levels of the drug in your body, which can make it less effective.



When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Tiredness and/or extreme weakness that interferes with your daily activities
- Wheezing and/or trouble breathing
- Coughing up yellow, green, or bloody mucus
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- No bowel movement in 3 days or when you feel uncomfortable
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Pain in your abdomen that does not go away
- Pain that does not go away, or is not relieved by prescribed medicines
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective non-hormonal methods of birth control during your cancer treatment and for 6 months after stopping treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 3 months after stopping treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- This medicine may interact with hormonal birth control (i.e., birth control pills, skin patches, vaginal rings, injections) and affect how they work. Discuss with your doctor and/or nurse what method of birth control may be right for you during your treatment.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 1 week after stopping treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised July 2023

