

Tepotinib (Tepmetko)

About This Drug

Tepotinib is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Nausea
- Diarrhea (loose bowel movements)
- Tiredness
- Swelling of the hands, feet, or any other part of the body
- Muscle and bone pain
- Trouble breathing

Note: Each of the side effects above was reported in 20% or greater of patients treated with tepotinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other drugs in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Inflammation (swelling) and/or scarring of the lungs, which can be life-threatening. You may have a dry cough or trouble breathing.
- Severe changes in your liver function, which can be life-threatening.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

How to Take Your Medication

- Swallow the medicine whole with food at approximately the same time each day. Do not crush, split, or chew tablets.
- If you have trouble swallowing, you can place the tablet(s) in a glass container containing 30 mL of non-carbonated water. Stir without crushing, until the tablet(s) is dispersed into small pieces (tablets will not completely dissolve) and drink immediately or within 1 hour. Do not chew pieces of

the tablet. After drinking, add an additional 30 mL of water to the container to rinse it, stir and drink right away.

- **Missed dose:** If you miss a dose, take it as soon as you think about it **ONLY** if your next dose is due in more than 8 hours. If your next dose is due in **LESS** than 8 hours, then skip the missed dose and take your next dose at the regular time. If you vomit a dose, take your next dose at the regular time. If you vomit a dose or miss a dose, contact your doctor.
- **Handling:** Wash your hands after handling your medicine. Caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- **Storage:** Store this medicine in the original container at room temperature.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise, with your doctor's approval. If you feel too tired to exercise vigorously, try taking a short walk.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- To help with nausea, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid). Eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop diarrhea.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.

Food and Drug Interactions

- There are no known interactions of tepotinib with food, however this medicine should be taken with food.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with tepotinib. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.



When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness or weakness that interferes with your daily activities
- Wheezing and/or trouble breathing
- Dry cough
- Pain in your chest
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Pain that does not go away, or is not relieved by prescribed medicines
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of hands, feet, or any other part of the body
- If you think you may be pregnant or may have impregnated your partner.

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 1 week after stopping treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for 1 week after stopping treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 1 week after stopping treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised April 2023

