

Vimseltinib (Romvimza)

About This Medicine

VIMSELTINIB (vim SEL ti nib) treats noncancerous tumors that occur in joints. It works by blocking a protein that causes tumor cells to grow and multiply. It is given orally (by mouth).

This medicine may be used for other purposes; ask your care team you have questions.

Possible Side Effects

- Decrease in the number of white blood cells. This may raise your risk of infection.
- Tiredness
- Swelling of your face and/or around your eyes
- Swelling of your hands, legs, ankles, and/or feet
- Increase in your cholesterol level
- Changes in your liver function
- Itching
- Rash

Note: Each of the side effects above was reported in 20% or greater of people treated with vimseltinib. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Severe changes in your liver function, which can cause liver failure and be life threatening.
- Tell your care team before taking this medicine if you have an allergy to FDandC Yellow No.5 (tartrazine) and No.6 (Sunset Yellow FCF) or aspirin. Vimseltinib contains FDandC Yellow No.5 which may cause allergic-type reactions in certain people, especially people who also have an allergy to aspirin.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

How to Take Your Medicine

- Swallow the medicine whole with or without food, 2 times a week and at least 72 hours apart. Do not break, chew, or open the capsules.
- **Missed dose:** If you miss a dose by less than 48 hours, take the missed dose right away and go back to your normal schedule. If you miss a dose by more than 48 hours, then skip the missed dose and go back to your normal schedule.
- If you vomit within 30 minutes of taking your dose, repeat the dose. If you vomit 30 minutes after taking your dose, do not take another dose and go back to your normal schedule.
- **Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
- **Handling:** Wash your hands with soap and water before and after handling your medicine. Your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the original container at room temperature. Keep out of the reach of children and pets.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medicine label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your care team how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your care team tells you, and whenever you feel like you may have a fever.
- If you get a rash do not put anything on it unless your care team says you may. Keep the area around the rash clean and dry. Ask your care team for medicine if your rash bothers you.
- To help with itching, moisturize your skin several times a day.
- Avoid sun exposure and apply sunscreen routinely when outdoors.

Food and Medicine Interactions

- There are no known interactions of vimseltinib with food .
- Check with your care team about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with vimseltinib. Also, check with your care team

before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling of the face and/or around your eyes
- Swelling of your hands, legs, ankles, and/or feet
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Itching that is bothersome
- A new rash or a rash that is not relieved by prescribed medicines

Reproduction Warnings

- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 1 month after the last dose. If you can get your partner pregnant, use birth control while taking this medicine and for 1 month after the last dose. Tell your care team right away if you think you might be pregnant or think your partner might be pregnant.
- **Breastfeeding warning:** Do not breastfeed while taking this medicine and for 1 month after the last dose.
- **Fertility warning:** This medicine may affect your ability to have children in the future. If you plan to have children, talk with your care team.

New February 2025

