

Zanidatamab-hrii (Ziihera)

About This Medicine

ZANIDATAMAB-HRII (ZAN i DAT a mab) treats bile duct cancer. It works by blocking a protein that causes cancer cells to grow and multiply. This helps to slow or stop the spread of cancer cells. It is a monoclonal antibody. It is given in the vein (IV).

This medicine may be used for other purposes; ask your care team if you have questions.

Possible Side Effects

- Diarrhea (loose bowel movements)
- Pain in your abdomen
- While you are getting this medicine in your vein (IV), you may have a reaction to the medicine. Sometimes you may be given medicine to stop or lessen these side effects. Your care team will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. These reactions may happen after your infusion. **If this happens, call 911 for emergency care.**
- Tiredness

Note: Each of the side effects above was reported in 20% or greater of people treated with zanidatamab-hrii. All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your care team.

Warnings and Precautions

- Changes in your heart's ability to pump blood properly

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your care team.

Important Information

- This medicine may be present in the urine, stool and other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine

or your body fluids. Follow safety precautions during your treatment and for as long as directed by your care team after your treatment. If you take a pill each day, follow these precautions every day.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Drink enough fluids to keep your urine pale yellow.
- If you have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- Eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping. Ask your care team about medicine that can lessen or stop diarrhea.
- Keeping your pain under control is important to your well-being. Please tell your doctor or care team if you are experiencing pain.
- Infusion reactions may occur after your infusion. **If this happens, call 911 for emergency care.**

Food and Medicine Interactions

- There are no known interactions of zanidatamab-hrii with food.
- This medicine may interact with other medicines. Tell your care team about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your care team before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call Your Care Team

Call your care team if you have any of these symptoms and/or any new or unusual symptoms:

- Tiredness that interferes with your daily activities
- Trouble breathing
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Pain in your abdomen that does not go away
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Swelling of the hands, feet, or any other part of the body
- Weight gain of 5 pounds in one week (fluid retention)
- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**

Reproduction Warnings



- **Pregnancy warning:** This medicine can cause serious birth defects. If you can become pregnant, use birth control while taking this medicine and for 4 months after the last dose. Tell your care team right away if you think you might be pregnant.
- **Breastfeeding warning:** Talk with your care team about breastfeeding during treatment. You may need to stop breastfeeding.
- **Fertility warning:** The effect of this medicine on fertility is not known. If you plan to have children, talk with your care team.

Revised January 2026

