

Client Name:

Date:

RDN/NDTR:

Email:

Phone:

Heart-Healthy Nutrition Therapy

A heart-healthy diet is recommended to reduce your unhealthy blood cholesterol levels, manage high blood pressure, and lower your risk for heart disease.

To follow a heart-healthy diet,

- Eat a balanced diet with whole grains, fruits and vegetables, and lean protein sources.
- Achieve and maintain a healthy weight.
- Choose heart-healthy unsaturated fats. Limit saturated fats, *trans* fats, and cholesterol intake.
- Eat more plant-based or vegetarian meals using beans and soy foods for protein.
- Eat whole, unprocessed foods to limit the amount of sodium (salt) you eat.
- Limit refined carbohydrates especially sugar, sweets and sugar-sweetened beverages.
- If you drink alcohol, do so in moderation: one serving per day (women) and two servings per day (men).
 - One serving is equivalent to 12 ounces beer, 5 ounces wine, or 1.5 ounces distilled spirits

Tips for Choosing Heart-Healthy Fats

Choose lean protein and low-fat dairy foods to reduce saturated fat intake.

- Saturated fat is usually found in animal-based protein and is associated with certain health risks. Saturated fat is the biggest contributor to raised low-density lipoprotein (LDL) cholesterol levels in the diet. Research shows that limiting saturated fat lowers unhealthy cholesterol levels. Eat no more than 5-6% of your total calories each day from saturated fat. Ask your registered dietitian nutritionist (RDN) to help you determine how much saturated fat is right for you.
- There are many foods that do not contain large amounts of saturated fats. Swapping these foods to replace foods high in saturated fats will help you limit the saturated fat you eat and improve your cholesterol levels. You can also try eating more plant-based or vegetarian meals.

Instead of...	Try:
▶ Whole milk, cheese, yogurt, and ice cream	▶ 1%, ½%, or skim milk, low-fat cheese, non-fat yogurt, and low-fat ice cream
▶ Fatty, marbled beef and pork	▶ Lean beef, pork, or venison
▶ Poultry with skin	▶ Poultry without skin
▶ Butter, stick margarine	▶ Reduced-fat, whipped, or liquid spreads
▶ Coconut oil, palm oil	▶ Liquid vegetable oils: corn, canola, olive, soybean and safflower oils

Avoid trans fats.

- *Trans* fats increase levels of LDL-cholesterol. Hydrogenated fat in processed foods is the main source of *trans* fats in foods.
- *Trans* fats can be found in stick margarine, shortening, processed sweets, baked goods, some fried foods, and packaged foods made with hydrogenated oils. Avoid foods with “partially hydrogenated oil” on the ingredient list such as: cookies, pastries, baked goods, biscuits, crackers, microwave popcorn, and frozen dinners.

Choose foods with heart healthy fats.

- Polyunsaturated and monounsaturated fat are unsaturated fats that may help lower your blood cholesterol level when used in place of saturated fat in your diet.
- Ask your RDN about taking a dietary supplement with plant sterols and stanols to help lower your cholesterol level.
- Substituting saturated fats with unsaturated fats is beneficial to cholesterol levels. Try these easy swaps:

Instead of...

Try:

■ Butter, stick margarine, or solid shortening	■ Reduced-fat, whipped, or liquid spreads
■ Beef, pork, or poultry with skin	■ Fish, seafood, and skinless poultry
■ Chips, crackers, snack foods	■ Raw or unsalted nuts and seeds or nut butters
	■ Hummus with vegetables
	■ Avocado on toast
■ Coconut oil, palm oil	■ Liquid vegetable oils: corn, canola, olive, soybean and safflower oils

Limit the amount of cholesterol you eat to less than 200 milligrams per day.

- Cholesterol is a substance carried through the bloodstream via lipoproteins, which are known as “transporters” of fat. Some body functions need cholesterol to work properly, but too much cholesterol in the bloodstream can damage arteries and build up blood vessel linings (which can lead to heart attack and stroke). You should eat less than 200 milligrams cholesterol per day.
- People respond differently to eating cholesterol. There is no test available right now that can figure out which people will respond more to dietary cholesterol and which will respond less. For individuals with high intake of dietary cholesterol, different types of increase (none, small, moderate, large) in LDL-cholesterol levels are all possible.
- Food sources of cholesterol include egg yolks and organ meats such as liver, gizzards. Limit egg yolks to two to four per week and avoid organ meats like liver and gizzards to control cholesterol intake.

Tips for Choosing Heart-Healthy Carbohydrates

Consume foods rich in viscous (soluble) fiber

- Viscous, or soluble, fiber is found in the walls of plant cells. Viscous fiber is found only in plant-based foods—animal-based foods like meat or dairy products do not contain fiber. In the stomach, viscous fibers absorb water and swell to form a thick, jelly-like mass. This helps to lower your unhealthy cholesterol.
 - Rich sources of viscous fiber include asparagus, Brussels sprouts, sweet potatoes, turnips, apricots, mangoes, oranges, legumes, barley, oats, and oat bran.
- Eat at least 5 to 10 grams of viscous fiber each day. As you increase your fiber intake gradually, also increase the amount of water you drink. This will help prevent constipation.
- If you have difficulty achieving this goal, ask your RDN about fiber laxatives. Choose fiber supplements made with viscous fibers such as psyllium seed husks or methylcellulose to help lower unhealthy cholesterol.

Limit refined carbohydrates

- There are three types of carbohydrates: starches, sugar, and fiber. Some carbohydrates occur naturally in food, like the starches in rice or corn or the sugars in fruits and milk. Refined carbohydrates—foods with high amounts of simple sugars—can raise triglyceride levels. High triglyceride levels are associated with coronary heart disease.
- Some examples of refined carbohydrate foods are table sugar, sweets, and beverages sweetened with added sugar.

Tips for Reducing Sodium (Salt)

Although sodium is important for your body to function, too much sodium can be harmful for people with high blood pressure. As sodium and fluid buildup in your tissues and bloodstream, your blood pressure increases. High blood pressure may cause damage to other organs and increase your risk for a stroke.

Keep your salt intake to 2300 milligrams or less per day. Even if you take a pill for blood pressure or a water pill (diuretic) to remove fluid, it is still important to have less salt in your diet. Ask your RDN what amount of sodium is right for you.

- Avoid processed foods. Eat more fresh foods.
 - Fresh fruits and vegetables are naturally low in sodium, as well as frozen vegetables and fruits that have no added juices or sauces.
 - Fresh meats are lower in sodium than processed meats, such as bacon, sausage, and hotdogs. Read the nutrition label or ask your butcher to help you find a fresh meat that is low in sodium.
- Eat less salt—at the table and when cooking.
 - A single teaspoon of table salt has 2,300 mg of sodium.
 - Leave the salt out of recipes for pasta, casseroles, and soups.
 - Ask your RDN how to cook your favorite recipes without sodium
- Be a smart shopper.
 - Look for food packages that say “salt-free” or “sodium-free.” These items contain less than 5 milligrams of sodium per serving.
 - “Very low-sodium” products contain less than 35 milligrams of sodium per serving.
 - “Low-sodium” products contain less than 140 milligrams of sodium per serving.
 - Beware of “reduced salt” or “reduced sodium” products. These items may still be high in sodium. Check the nutrition label.
- Add flavors to your food without adding sodium.
 - Try lemon juice, lime juice, fruit juice or vinegar.
 - Dry or fresh herbs add flavor. Try basil, bay leaf, dill, rosemary, parsley, sage, dry mustard, nutmeg, thyme, and paprika.
 - Pepper, red pepper flakes, and cayenne pepper can add spice to your meals without adding sodium. Hot sauce contains sodium, but if you use just a drop or two, it will not add up to much.
- Buy a sodium-free seasoning blend or make your own at home.

Additional Lifestyle Tips

Achieve and maintain a healthy weight.

- Talk with your RDN or your doctor about what is a healthy weight for you.
- Set goals to reach and maintain that weight.
- To lose weight, reduce your calorie intake along with increasing your physical activity. A weight loss of 10 to 15 pounds could reduce LDL-cholesterol by 5 milligrams per deciliter.

Participate in physical activity.

- Talk with your health care team to find out what types of physical activity are best for you. Set a plan to get about 30 minutes of exercise on most days.

Foods to Choose or to Limit

Food Group	Foods to Choose	Foods to Limit
Grains	<ul style="list-style-type: none"> ▶ Whole grain breads and cereals, including whole wheat, barley, rye, buckwheat, corn, teff, quinoa, millet, amaranth, brown or wild rice, sorghum, and oats ▶ Pasta, especially whole wheat or other whole grain types ▶ Brown rice, quinoa or wild rice ▶ Whole grain crackers, bread, rolls, pitas ▶ Home-made bread with reduced-sodium baking soda 	<ul style="list-style-type: none"> ▶ Breads or crackers topped with salt ▶ Cereals (hot or cold) with more than 300 mg sodium per serving ▶ Biscuits, cornbread, and other “quick” breads prepared with baking soda ▶ Bread crumbs or stuffing mix from a store ▶ High-fat bakery products, such as doughnuts, biscuits, croissants, danish pastries, pies, cookies ▶ Instant cooking foods to which you add hot water and stir—potatoes, noodles, rice, etc. ▶ Packaged starchy foods—seasoned noodle or rice dishes, stuffing mix, macaroni and cheese dinner ▶ Snacks made with partially hydrogenated oils, including chips, cheese puffs, snack mixes, regular crackers, butter-flavored popcorn
Protein Foods	<ul style="list-style-type: none"> ▶ Lean cuts of beef and pork (loin, leg, round, extra lean hamburger) ▶ Skinless poultry ▶ Fish ▶ Venison and other wild game ▶ Dried beans and peas ▶ Nuts and nut butters (unsalted) ▶ Seeds and seed butters (unsalted) ▶ Meat alternatives made with soy or textured vegetable protein ▶ Egg whites or egg substitute ▶ Cold cuts made with lean meat or soy protein 	<ul style="list-style-type: none"> ▶ Higher-fat cuts of meats (ribs, t-bone steak, regular hamburger) ▶ Bacon, sausage, or hot dogs ▶ Cold cuts, such as salami or bologna, deli meats, cured meats, corned beef ▶ Organ meats (liver, brains, gizzards, sweetbreads) ▶ Poultry with skin ▶ Fried or smoked meat, poultry, and fish ▶ Whole eggs and egg yolks (more than 2-4 per week) ▶ Salted legumes, nuts, seeds, or nut/seed butters ▶ Meat alternatives with high levels of sodium (>300 mg per serving) or saturated fat (>5 g per serving)
Dairy and Dairy Alternatives	<ul style="list-style-type: none"> ▶ Nonfat (skim), low-fat, or 1%-fat milk ▶ Nonfat or low-fat yogurt or cottage cheese ▶ Fat-free and low-fat cheese ▶ Fortified non-dairy milk: almond, cashew, pea, and soy 	<ul style="list-style-type: none"> ▶ Whole milk, 2% fat milk, buttermilk ▶ Whole milk yogurt or ice cream ▶ Cream, half-and-half ▶ Cream cheese ▶ Sour cream ▶ Cheese

Food Group	Foods to Choose	Foods to Limit
Vegetables	<ul style="list-style-type: none"> ▶ Fresh, frozen, or canned vegetables without added fat or salt ▶ Avocados 	<ul style="list-style-type: none"> ▶ Canned or frozen vegetables with salt, fresh vegetables prepared with salt, butter, cheese, or cream sauce ▶ Fried vegetables ▶ Pickled vegetables such as olives, pickles, or sauerkraut
Fruits	<ul style="list-style-type: none"> ▶ Fresh, frozen, canned, or dried fruit 	<ul style="list-style-type: none"> ▶ Fried fruits ▶ Fruits served with butter or cream
Oils	<ul style="list-style-type: none"> ▶ Unsaturated oils (corn, olive, peanut, soy, sunflower, canola) ▶ Soft or liquid margarines and vegetable oil spreads ▶ Salad dressings made from unsaturated fats 	<ul style="list-style-type: none"> ▶ Butter, stick margarine, shortening ▶ Partially hydrogenated oils or trans fats ▶ Tropical oils (coconut, palm, palm kernel oils)
Other	<ul style="list-style-type: none"> ▶ Prepared or homemade foods, including soups, casseroles, and salads made from recommended ingredients and contain <600 mg sodium ▶ Low-sodium seasonings (ketchup, barbeque sauce) ▶ Spices, herbs, salt-free seasoning mixes and marinades ▶ Vinegar ▶ Lemon or lime juice 	<ul style="list-style-type: none"> ▶ Prepared foods, including soups, casseroles, and salads made from recommended ingredients and contain >600 mg sodium ▶ Frozen meals and prepared sides that are >600 mg of sodium ▶ Sugary and/or fatty desserts, candy, and other sweets ▶ Salts: sea salt, kosher salt, onion salt, and garlic salt, seasoning mixes containing salt ▶ Flavorings: bouillon cubes, catsup or ketchup, barbeque sauce, Worcestershire sauce, soy sauce, salsa, relish, teriyaki sauce

Heart-Healthy Sample 1-Day Menu

Meal	Menu
Breakfast	<ul style="list-style-type: none">■ 1 cup oatmeal■ 1 cup blueberries■ 1 ounce almonds, unsalted■ 1 cup skim milk■ 1 cup brewed coffee
Lunch	<ul style="list-style-type: none">■ 2 slices whole wheat bread■ 2 ounces lean turkey breast■ 1 ounce low-fat Swiss cheese■ 1 teaspoon mustard■ 1 sliced tomato■ 1 lettuce leaf■ 1 pear■ 1 cup skim milk
Afternoon Snack	<ul style="list-style-type: none">■ 1 ounce trail mix with unsalted nuts, seeds, and raisins
Dinner	<ul style="list-style-type: none">■ 3 ounces broiled salmon■ ⅔ cup brown rice■ 1 teaspoon margarine, soft tub■ ½ cup broccoli, cooked■ ½ cup carrots, cooked■ 1 cup tossed salad■ 1 teaspoon olive oil and vinegar dressing■ 1 small whole grain roll■ 1 teaspoon margarine, soft tub■ 1 cup tea
Evening Snack	<ul style="list-style-type: none">■ 1 small banana

Sample Meal Plan

Use this form to create an individualized meal plan.

Meal	Menu
Breakfast	
Lunch	
Dinner	
Snack	

Notes: