



Breastfeeding Guide

HOW TO GET OFF
TO A GREAT START



Breastfeeding Guide - How to Get Off to a Great Start

Everyone needs good nutrition. But it is even more important for babies and children because they need good nutrition in order to grow and develop.

Your breastmilk is more than nutrition. It also protects your baby. Your breastmilk helps keep your baby from getting sick. And it lowers your baby's risk of asthma, allergies, and sudden infant death syndrome (SIDS).

Your breastmilk protects your child for a long time, long after you stop breastfeeding. For example, it lowers your baby's risk of being overweight later in life. And it lowers his risk for adult diabetes and some types of cancer.

Breastfeeding is good for you, too. It lowers your risk of breast cancer, ovarian cancer and diabetes. And it burns about 600 calories a day, making it easier to lose some of the weight you gained while you were pregnant. Breastfeeding also helps in other ways. It helps your baby's brain develop. It creates a close bond between you and your baby. It saves money in formula and health care costs and cuts down on the days you miss from work. And breastmilk is a natural resource, so it helps our planet.

By breastfeeding, you are doing something that is good for your health and your baby's health - for life.

Congratulations for breastfeeding!






table of content



BREASTFEEDING GUIDE - HOW TO GET OFF TO A GREAT START

- 2 DURING PREGNANCY
Your body is built for breastfeeding
How long should I breastfeed?
- 3 FIRST HOUR
Tips for getting breastfeeding off to a good start
Your baby's first hour
- 4 COLOSTRUM
Hand expression of colostrum
Once you are in your room
- 5 FEEDING POSITIONS
How do I hold my baby for breastfeeding?
- 6 COMFORTABLE LATCH
How do I attach my baby to my breast?
- 7 SIGNS OF HUNGER AND FULL
How often should I feed my baby?
How do I tell if my baby is full?
- 8 WAKE YOUR BABY
Sleepy babies
- 9 GAINING WEIGHT
How do I tell if my baby is getting enough to eat?
- 10 REDUCING THE SWELLING
My breasts feel too full. Do I have too much milk?
Do I have too little milk?
- 11 TRACKING
First Week Daily Breastfeeding Log
How do I know if breastfeeding is going well?
- 13 TIPS
How do I know when to call the doctor?
How do I take care of myself?



Your body is built for breastfeeding

During pregnancy, your body gets ready to breastfeed. In the fourth month of pregnancy, your breasts start making colostrum, the first milk for your baby. Colostrum can be yellow or clear. The colostrum that you feed your baby during the first few days after birth will boost his immune system and help him poop. This lowers his chance of jaundice, a condition where the baby's skin and eyes turn yellow.

Your newborn baby has a tiny stomach – about the size of a small marble. On the first day, his stomach can only hold about one teaspoon of colostrum each time you feed him. Your baby's stomach and appetite will grow as he grows.


Colostrum is the only thing your baby needs the first few days after he is born.



How long should I breastfeed?


Any amount of breastfeeding is good, but the longer you breastfeed, the better. The American Academy of Pediatrics says that women should breastfeed their babies “exclusively” (feed only breastmilk to their babies) for about the first 6 months. At that point, mothers should start giving baby foods and continue to breastfeed, at least until the baby is 12 months old. Experts say that women should keep breastfeeding after one year as long as the mom and the baby both want to.

Tips for getting breastfeeding off to a good start



These steps can help you breastfeed. Let your nurse know that you want to follow these steps.

1. Breastfeed within the first hour after birth, even if you had a caesarean (or c-section) birth.
2. Feed only breastmilk to your baby. Ask for a breast pump if your baby is not feeding from your breast.
3. Hold your baby “skin-to-skin” as much as you can. This means that your baby’s skin is touching your skin.
4. Keep your baby in your room, day and night, except for special procedures.
5. Do not give a pacifier or bottle to your baby.



Your baby’s first hour



When your baby is born, he should be placed on your bare chest. This skin-to-skin contact will comfort your baby after the delivery. The staff can do most things they need to do while you are holding your baby. Your baby will start to look for your breast and nipple. His head will bob up and down as he crawls to your breast. It may take some time, but your baby will attach to your breast when he is ready. It is truly amazing to watch your baby find your breast and start feeding.

If he does not show interest in the first hour, place his cheek on top of one of your breasts so he can feel, smell and taste your nipple. If he still does not show interest in eating, ask for help. Your baby should breastfeed within the first hour or so after birth.

Hand expression of colostrum

Some babies are extra sleepy in the first few days or need more time to practice breastfeeding. If your baby will not feed from your breast, you may need to remove your breastmilk by hand. This is called “hand expression.” Hand expression of colostrum is better than pumping because colostrum can stick to pump parts, so you might not collect as much of it.

To hand express:

1. Wash your hands.
2. Use a clean container with a wide opening or a spoon to collect your colostrum.
3. Hold the container near your nipple. With your other hand, place your fingers and thumb in line with your nipple and about one to two inches away from your nipple.
 - * Press your finger and thumb toward your ribs.
 - * Squeeze gently and catch your milk in the cup.
 - * Relax your hand.
 - * Repeat the process. Press in, squeeze gently, relax, and repeat.
4. It may take a few minutes before you see any colostrum. Your baby only needs about one teaspoon each time you feed him.
5. Ask a nurse or lactation consultant to help you express and feed your colostrum to your baby.




Once you are in your room

- Ask to keep your baby in your room at all times – day and night.
- Feed only breastmilk to your baby. Do not give bottles or pacifiers.
- Hold your baby skin-to-skin as much as possible.

Babies love skin-to-skin contact. Once you are in your room, dress your baby in just a diaper, cap, and socks and hold her against your bare chest. Cover her back with a blanket. Tuck the ends of the blanket behind your own back to keep her snuggled up close to you. You can also put a blanket around your shoulders, too. Skin-to-skin contact with dad is also a great way for dads to bond with their babies.

Holding babies skin-to-skin helps them:

- Breastfeed better
- Cry less
- Recover from the delivery more quickly



How do I hold my baby for breastfeeding?

There are many positions you can use. Try different positions to find what works best for you.



Laid Back Hold

- Lay back and use pillows for support and comfort.
- Place your baby face down between your breasts.
- Let your baby search, crawl up, nuzzle, and attach to your breast. Help your baby in whatever way feels natural.
- Place a blanket across your baby's back, if needed.



Cross-Cradle Hold

- Place a pillow in your lap to bring your baby up to breast level. Put your baby on the pillow, tummy to tummy with you, with the baby's nose across from your nipple.
- Support your baby's head by holding your hand at the base of his skull.
- Lift your breast to bring your nipple up to your baby's nose. To lift your breast, lean back and place your fingers below your breast near your ribs, keeping your hand far away from the nipple.
- Once your baby is latched on, you can let go of your breast and bring your arm around your baby into a regular cradle-hold. If your breast are large, you may need to support your breast the whole time.



Clutch (Football) Hold

- Place a pillow at your side.
- Put your baby on the pillow with his legs under your arm.
- Slide your arm under your baby's back. Support the base of his head and neck with your hand.



Side-Lying Hold

- Lie on your side with knees bent. Place pillows between your knees, under your head and neck, and behind your back, if needed.
- Put your baby on his side, facing your nipple.
- Support your baby by placing your arm, a pillow, or a rolled-up blanket behind him.





How do I attach my baby to my breast?

If you are using the laid back position, allow your baby to latch on when he is ready. If you are using a different position, you will control the attachment more than your baby.



1

Follow these tips

Hold your baby so that his nose is in line with your nipple. Touch your baby's nose and upper lip with your nipple.



2

Wait until his mouth opens very wide.



3

Quickly bring the baby onto your nipple and breast so that his chin touches your breast first and he gets a large mouthful of nipple and breast.



4

If you feel painful tugging or pinching, slide your finger into the corner of his mouth to break the suction and try again. It may take a few tries to get a good, comfortable latch.

You can tell your baby is attached well if:

- You are not feeling sharp pain. Gentle tugging is normal.
- Both of your baby's lips are flipped out, not pulled in.
- More of the bottom of your areola (the dark area around nipple) is in the baby's mouth than the top.
- His chin is buried in your breast with his nose tipped away slightly or lightly touching.
- His mouth is as wide open as a yawn.

After the first two days, you may also:

- Hear your baby swallow; or
- See milk leaking from your baby's mouth or your other breast.

Your nipple should look the same coming out of your baby's mouth as it did going in. If your nipple looks pinched when it first comes out of your baby's mouth, your baby is not attaching well. About half of all babies don't attach well on the first day. Ask for help right away and be patient. It may take a few days for you and your baby to learn the art of breastfeeding.

How often should I feed my baby?

Your baby is ready to feed any time he shows **early signs of hunger**, which are:

- Eye movements under eyelids
- Sucking motions while sleeping
- Restless body movements while sleeping
- Bringing hands toward mouth

Don't wait until he is fussy or crying. That makes it harder to breastfeed.

If someone offers to take your baby to the nursery so you can rest, tell them you would like to keep your baby in your room. Room-sharing protects you and your baby from infection, and helps you both sleep better. Also, you will get more skin-to-skin contact with your baby, and you will learn your baby's hunger cues faster.




How do I tell if my baby is full?

Your baby will signal when he is full when he:

- Lets go of your breast and nipple
- Falls asleep and stops sucking
- Relaxes his hands and body.

If your baby comes off the breast relaxed and sleepy he has probably had a good feeding. If he stops sucking and does not come off the breast on his own, slide your finger into the corner of his mouth to break the suction. Burp him and offer the other breast. He may nurse again right away or he may take the other side in a few minutes or a few hours.

If your baby comes off the breast crying, he may not be getting a good latch. Ask for help right away.



Sleepy babies

Sleepy babies are not good at showing signs that they are hungry.

Try waking your baby first with a variety of motions and sounds, such as different positions, using different words and sounds, and touching different parts of his body. Then get ready to feed.


- Place your baby in skin-to-skin contact and gently rub his back, arms, hands, and feet.
- Talk to him in a calming voice.
- When he begins to move more or opens his eyes, move him to your breast to feed.
- When he stops sucking, gently squeeze and massage your breast. Move your hand to a different area of your breast to massage and squeeze every time your baby stops sucking.

Most sleepy babies will get better at showing hunger cues in a few days. It may take 10 to 15 minutes to wake a sleepy baby.

Sleepy babies should:

- be encouraged to nurse at least 8 times a day or every 1–3 hours during the day, and
- have no more than one four-hour sleep period in 24 hours.





How do I tell if my baby is getting enough to eat?

Weight gain is the best way to tell if your baby is getting enough to eat. Tracking how many wet and poopy diapers your baby has can also be helpful.

Your baby may have some pale yellow, pink or red wet diapers and some black, brown or greenish poopy diapers in the first two to three days.

Baby's Average Diaper Output in the First 4 Weeks				
Day	Wet Diapers	Urine Color	Poopy Diapers	Poop Color
1	Number will vary	Yellow/Pink/Red	Number will vary	Black
2	Number will vary	Yellow/Pink/Red	Number will vary	Blackish/Green
3	3 or more	Yellow/Pink/Red	3 or more	Greenish/Yellow
4	4 or more	Clear/Pale Yellow	3 or more	Greenish/Yellow, Seedy
5 to 28	6 or more	Clear/Pale Yellow	3 or more	Yellow, Seedy

Your baby's poop can look watery and seedy and at other times it will look more like cottage cheese. By day six, many breastfed babies will poop in almost every diaper. At least three or four of those diapers should have a large amount of poop - about the size of a golf ball, but soft and spread out. After 4-6 weeks, a baby may only poop a few times a week.



Day 1





Day 3-4



Day 5

Newborns often lose weight in the first couple days of life. Your baby should regain his birth weight by day 10-14. After he regains his birth weight, he should gain about 4 to 8 ounces a week. Be sure to see your baby's health care provider within 3-5 days of hospital discharge to make sure your baby is gaining weight. Use the **First Week Daily Breastfeeding Log** on page 11 of this book to track your baby's diapers. When your baby is four days old, complete the **How do I know if breastfeeding is going well survey**, on page 12.






My breasts feel too full. Do I have too much milk?

When your baby is 2 to 5 days old, your milk supply will increase and your breasts will feel heavier. Nurse your baby often to keep your breasts from becoming too full.

If your breasts get too full, take a short, warm shower or put a warm, wet towel over your breasts and nipples for a minute or two and hand express a little milk before each feeding. Gently massage your breasts toward your nipples while your baby is breastfeeding. Cold packs can be used for up to 20 minutes between feedings if the fullness becomes uncomfortable.

If your breasts become so full that you have trouble hand expressing, try the softening method found at <http://www.breastmilkcounts.com/educational-activities.html>.



Do I have too little milk?


Many babies will nurse very often at one time of the day. Many babies do this in the evening. This is normal. The more often you nurse your baby, the more milk you will make. Breastfeed often and do not give your baby formula. Exclusive breastfeeding for the first few weeks is very important for building a good milk supply.

Sometime between your baby's first and third week of life, he may want to nurse more often. About this same time your breasts will naturally soften and feel less full. This does not mean you don't have enough milk. It means your baby is having his first growth spurt and your milk supply is changing to meet your baby's needs.

Try not to give your baby formula. Instead, set up a little area or "nest" by gathering plenty of supplies around you such as snacks, drinks, phone, remote controls, books, and magazines. Try to spend as much time as you can in your "nest" with your baby so you can get to know and enjoy your baby. Relax and nurse as often as your baby wants.

Growth spurts usually happen around these times.

- 1 to 3 weeks of age
- 6 weeks of age
- 3 months of age



First Week Daily Breastfeeding Log

Baby's Name:

Baby's Birth Date:

Birth Time:

Birthweight:

- Fill in the start time and day of week in each box. The start time should be the time your baby was born and should be the same for each day.
- The day of week should change each day.
- Circle a B each time your baby breastfeeds.
- Circle a W each time your baby has a wet diaper.
- Circle an S each time your baby has a soiled or dirty diaper.
- Circle the plus sign if your baby has more than the daily goal for wet and soiled diapers.

Day 1 (1st 24 hours)

Start Time:	Day of Week:										Goal	
Breastfeedings	B	B	B	B	B	B	B	B	B	B	+	6+
Wet diapers	(pale yellow/pink/red)											Any
Soiled diapers	(black tarry)											Any

Day 2 (2nd 24 hours)

Start Time:	Day of Week:										Goal	
Breastfeedings	B	B	B	B	B	B	B	B	B	B	+	8-12+
Wet diapers	(pale yellow/pink/red)											Any
Soiled diapers	(black tarry)											Any

Day 3 (3rd 24 hours)

Start Time:	Day of Week:										Goal		
Breastfeedings	B	B	B	B	B	B	B	B	B	B	+	8-12+	
Wet diapers	W	W	+	(pale yellow/pink/red)									3-5+
Soiled diapers	S	S	S	+	(greenish)							3-5+	

Day 4 (4th 24 hours)

Start Time:	Day of Week:										Goal	
Breastfeedings	B	B	B	B	B	B	B	B	B	B	+	8-12+
Wet diapers	W	W	W	+	(clear/pale yellow)							3-5+
Soiled diapers	S	S	S	S	+	(greenish to yellow)						3-4+

Day 5 (5th 24 hours)

Start Time:	Day of Week:										Goal	
Breastfeedings	B	B	B	B	B	B	B	B	B	B	+	8-12+
Wet diapers	W	W	W	W	+	(clear/pale yellow)						4-6+
Soiled diapers	S	S	S	S	+	(yellow, seedy)						3-4+

Day 6-28

Start Time:	Day of Week:										Goal	
Breastfeedings	B	B	B	B	B	B	B	B	B	B	+	8-12+
Wet diapers	W	W	W	W	W	+	(clear/pale yellow)					4-6+
Soiled diapers	S	S	S	S	+	(yellow, seedy)						3-4+

Let your baby breastfeed until he ends the feeding. Burp him and offer the other side each time. He may fill up on one breast but you should always offer the other to be sure. You CAN'T nurse too often. You CAN nurse too little.

A breastfed baby's poop is normally loose and unformed and will change colors in the first several days. It should be yellow by day four. It may appear seedy to cottage cheesy.

Your baby is probably getting enough to eat in his first week if he is meeting the goals for the number of daily breastfeedings and wet and dirty diapers and regains his birth weight by day 10-14. Some babies will go for days without a poop and this is normal too.

If you have question about how to use this log, please call your WIC breastfeeding counselor. If your baby is not meeting the goals, call your doctor and your WIC breastfeeding counselor.

Take this log to your baby's doctor at his first office visit which should be at 3-5 days of age.



How do I know if breastfeeding is going well?

The following questions will help you know whether you and your baby are off to a good start with breastfeeding.

Answer these questions when your baby is 4 days old.

Do you feel breastfeeding is going well?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Are your baby's swallows easier to hear since his first days in the hospital?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Is your baby able to latch on to your breasts without causing you discomfort?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Does your baby let you know when he/she is hungry?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Is your baby alert and active? (Answer no if you have to wake your baby for most feedings.)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Does your baby end each feeding by coming off your breast on his own?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Does your baby seem calm and satisfied after feedings?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Do your breasts feel full before feedings?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Do you have any sore, tender areas in your breast that are firm and red?	<input type="checkbox"/> No	<input type="checkbox"/> Yes
Does your baby nurse at least 8-12 times a day?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Is your baby waking at least one time during the night to breastfeed?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Are your nipples sore or do they look pinched when your baby comes off your breast?	<input type="checkbox"/> No	<input type="checkbox"/> Yes
Is your baby having bowel movements that are soft and mustard-yellow?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Is your baby having 3 or more poopy diapers each day that are larger than the size of a quarter?	<input type="checkbox"/> Yes	<input type="checkbox"/> No
When you notice a wet diaper, it is clear to pale yellow? (Pink to red wet diapers after day four can be a sign of dehydration.)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Is your baby having at least 3-5 wet diapers in 24 hours?	<input type="checkbox"/> Yes	<input type="checkbox"/> No

If you only checked answers in the left column, you are doing well!



If you checked any answers in the right column, call your WIC breastfeeding counselor right away.


Call your WIC breastfeeding counselor any time you have any questions or concerns about breastfeeding.



How do I know when to call the doctor?



You should call your baby's doctor if he:



- Does not regain his birth weight by 2 weeks of age.
- Has fewer than six wet diapers a day by day six.
- Has fewer than three poops a day by day three.
- Still has black poop on day four.
- Will not wake up to nurse at least eight times a day.
- Falls asleep or stops nursing right after attaching to your breast.

How do I take care of myself?

Forget about housework and try to sleep when your baby sleeps. When you are tired, lie down for feedings. Have snacks and drinks beside you each time you sit down to nurse. Keep meals simple — like a sandwich, soup, and fruit. Limit drinks with caffeine to no more than three a day. Most medicines are safe to take when you are breastfeeding but check with your doctor to make sure. Or call the Infant Risk Center at 1-806-352-2519.

Many women have mildly sore nipples the first few days of breastfeeding. If the discomfort only happens at the beginning of feedings and goes away when your milk starts to flow or within 30 seconds, this is probably normal. To soothe sore nipples, rub breastmilk into them or use a lanolin cream that is made for breastfeeding mothers.

If the soreness lasts the entire feeding or is severe, see a breastfeeding counselor right away. If you have soreness at the beginning of feeding only and it does not go away within two weeks, see a breastfeeding counselor. Call your local WIC clinic for breastfeeding help. Most likely, you are sore because of the way the baby is positioned or the way the baby is latching on. The soreness should go away after someone helps you figure out the cause.

Hang in there. Breastfeeding gets easier. Take it one day at a time and be proud of what you are doing for your baby. You are giving your baby the best possible start in life.





BREASTFEEDING HELP AND INFORMATION

Would you like to talk to someone about breastfeeding?

Most WIC clinics have breastfeeding Peer Counselors who can help with basic breastfeeding issues and International Board Certified Lactation Consultants to help with more difficult problems. Peer Counselors are WIC moms just like you who have breastfed their own children. WIC trains them to help other moms. Call your WIC clinic to speak to a breastfeeding Peer Counselor.



Visit the Lactation Care Center, the Lactation Foundation or Mom's Place on Facebook to ask a question or call to talk to a Lactation Consultant.

Mom's Place, Austin - 1-800-514-6667

The Lactation Foundation, Houston - 1-877-550-5008

The Lactation Care Center, Dallas – 1-855-550-6667



Join an online mom-to-mom breastfeeding group.

<http://forums.llli.org/>

<http://www.breastfeeding.com/forums/>

<http://messageboards.ivillage.com/ivpsbreastfeed?ice=ivl,searchmb>

<http://forum.kellymom.net/>


**For more breastfeeding information,
visit www.breastmilkcounts.com.**




Mother's Milk or Formula?

How you feed your baby may or may not be an easy choice for you but it is an important one.

Your baby's growth and health are determined by "building blocks" that are in the foods your baby receives.



Mother's Milk



Formula




Formula increases health risks for moms and babies.

Baby:

- Fussiness
- Constipation
- Colic
- Vomiting
- Allergies
- Asthma
- Ear infections
- Cancer
- Diabetes
- Respiratory infections
- Obesity
- SIDS
- Lower IQ

Mom:

- Breast cancer
- Uterine cancer
- Osteoporosis
- Anemia
- Obesity



Nursing your baby for even a few weeks is better than not at all. It's best to keep nursing. The longer you do, the more protection you and your baby receive.

Mother's milk *only* for the first 6 months
Mother's milk plus baby foods after 6 months



Breastfeeding Awareness and Support Program
 Healthy Baby Line
 1-800-986-BABY



Pennsylvania WIC Program
 1-800-WIC-WINS
 H511.940P June 2009

It's my birthday, *give me a hug!*

Skin-to-skin contact for you and your baby

What's "skin-to-skin"?

Skin-to-skin means your baby is placed belly-down, directly on your chest, right after she is born. Your care provider dries her off, puts on a hat, and covers her with a warm blanket, and gets her settled on your chest. The first hours of snuggling skin-to-skin let you and your baby get to know each other. They also have important health benefits. If she needs to meet the pediatricians first, or if you deliver by c-section, you can unwrap her and cuddle shortly after birth. Newborns crave skin-to-skin contact, but it's sometimes overwhelming for new moms. It's ok to start slowly as you get to know your baby.

Breastfeeding

Snuggling gives you and your baby the best start for breastfeeding. Eight different research studies have shown that skin-to-skin babies breastfeed better. They also keep nursing an average of six weeks longer. The American Academy of Pediatrics recommends that all breastfeeding babies spend time skin-to-skin right after birth. Keeping your baby skin-to-skin in his first few weeks makes it easy to know when to feed him, especially if he is a little sleepy.

A smooth transition

Your chest is the best place for your baby to adjust to life in the outside world. Compared with babies who are swaddled or kept in a crib, skin-to-skin babies stay warmer and calmer, cry less, and have better blood sugars.

Bonding

Skin-to-skin cuddling may affect how you relate with your baby. Researchers have watched mothers and infants in the first few days after birth, and they noticed that skin-to-skin moms touch and cuddle their babies more. Even a year later, skin-to-skin moms snuggled more with their babies during a visit to their pediatrician.



Photo © 2005 Pascale Wowak

Skin-to-skin beyond the delivery room

Keep cuddling skin-to-skin after you leave the hospital—your baby will stay warm and comfortable on your chest, and the benefits for bonding, soothing, and breastfeeding likely continue well after birth. Skin-to-skin can help keep your baby interested in nursing if he's sleepy. Dads can snuggle, too. Fathers and mothers who hold babies skin-to-skin help keep them calm and cozy.

About the research

Multiple studies over the past 30 years have shown the benefits of skin-to-skin contact. For more information, see Moore, ER, Anderson, GC, Bergman N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev.* 2007 Jul 18;(3):CD003519



**Massachusetts
Breastfeeding
Coalition**

254 Conant Road, Weston, MA 02493
www.massbreastfeeding.org | © 2011 MBC and Alison Stuebe



It's only natural

mother's love. mother's milk.



Cross Cradle Hold



Football Hold



Side-lying



Cradle Hold



Breastfeeding takes patience and practice at first. With the right support and information you can make it work.

Simple steps to breastfeeding success

Position yourself comfortably

- Sit up in bed, in a comfortable chair, or in a rocking chair (unless you are side-lying).
- You can use pillows to support your back and arms while you hold your baby.
- Relax your muscles and get comfortable.

Position your baby

- Hold your baby lying on his/her side with his/her body facing you.
- Hold your baby at the level of your nipple so that you're not leaning forward. Using the pillow on your lap will help.
- Use your hand to form a "U" around the darker part of your breast (areola) and offer your breast to the baby. Your thumb should be in line with the baby's nose, with the rest of your fingers on the other side of your breast.
- Your baby's head should be tilted back slightly to make it easy to suck and swallow.
- As shown here, there are several types of holding positions you can try to find the one that is right for you.
- Remember that making sure both you and your baby are comfortable is important to breastfeeding success.
- You can find more information about positioning and feeding your baby at www.womenshealth.gov/ItsOnlyNatural

Help your baby to latch properly

- Tickle the lips so your baby opens wide. Some women feel a pinch at first. However, if it lasts longer, your baby may not be latched correctly.
- Pull your baby close so the chin and lower jaw move into your breast first.
- Put the lower lip as far from the base of the nipple as possible, so your baby takes a large mouthful of breast.
- Remember your baby can breathe while at your breast because nostrils open wider to let air in.
- Some babies latch on right away, and some take more time.

Signs of a good latch

- The latch feels comfortable to you.
- Your baby's chest is against your body so the head doesn't need to turn.
- Your baby's mouth is filled with breast.
- You hear or see your baby swallow. For some babies the only sign of swallowing may be a pause in breathing.
- Your baby's ears "wiggle" slightly.
- Your baby's lips turn out, not in.



About It's Only Natural

Breastfeeding provides mothers and their babies with a healthy start. The U.S. Department of Health and Human Services created *It's Only Natural* to offer African American moms the knowledge, help, and support they need to breastfeed. You'll find all this at www.womenshealth.gov/ItsOnlyNatural. Breastfeeding. It's only natural with mother's love, mother's milk.



U.S. Department of Health and Human Services

How to know if your baby is getting enough milk

You may be concerned that you're not making enough milk. But your baby is likely getting more than you think at each feeding. Regardless of your baby's weight, a newborn's stomach is only the size of an almond.



Many babies lose a small amount of weight after birth, so don't be alarmed. Your baby's doctor will check weight at the first visit after you leave the hospital, which should be scheduled 2 to 3 days after birth. After the first few days, typical weight gain is a little less than 1 ounce per day. If he/she is getting enough milk, your baby will:

- Pass enough clear or pale yellow urine. It should not be deep yellow or orange. In the early days, your baby should have at least 1 wet diaper for each day of life (1 on day 1, 2 on day 2, etc.). Once the amount of milk you are making increases (24-72 hours), expect at least, 5-6 wet diapers every 24 hours.
- Have enough bowel movements. In the early days, your baby should have at least 1 dirty diaper for each day of life (1 on day 1, 2 on day 2, etc.). After day 4, expect at least 3-4 stools daily.
- Switch between short sleeping periods and wakeful, alert periods.
- Be satisfied and content right after feedings. (Keep in mind that breast milk digests quickly so babies must eat frequently.)
- Also, your breasts may feel less full after feeding your baby.

Talk to your baby's doctor or a breastfeeding expert if you are concerned that your baby is not getting enough milk before you stop breastfeeding or begin using formula.

 **It's only
natural**
mother's love. mother's milk.

Information for breastfeeding families

Hand Expression



Hand expression is a handy skill to have whenever you need to empty your breasts and you are not with your baby or your baby is temporarily unable to breastfeed. In the first few days after birth hand expression can be more effective at removing colostrum than using a breast pump. If your baby needs a supplement in the first few, use hand expression to provide the milk he needs!

Hand expression routine:

1. Apply heat, massage and stroke breasts
2. Position fingers behind areola
3. Press back towards chest
4. Compress fingers together to express milk
5. Relax and repeat getting a rhythm going
6. Express for 5-7 minutes
7. Move fingers to a different position
8. Massage and stroke the breast
9. Press back towards chest
10. Compress fingers together to express milk
11. Express milk for 3-5 minutes
12. Massage and stroke breasts
13. Move fingers to a different position
14. Express milk for 1-2 minutes
15. Complete cycle takes 20-30 minutes



Watch this video while you are hand expressing to see the technique in action!


<http://newborns.stanford.edu/Breastfeeding/HandExpression.html>

Provided for you by Lactation Education Resources. May be freely copied and distributed. Please be aware that the information provided is intended solely for general educational and informational purposes only. It is neither intended nor implied to be a substitute for professional medical advice. Always seek the advice of your physician for any questions you may have regarding your or your infant's medical condition. Never disregard professional medical advice or delay in seeking it because of something you have received in this information.

Breastfeeding

HAND EXPRESSION

Many women find that hand expression is an efficient way to pump when only occasional expression is necessary. In the few first days, when colostrum (first milk) is present and milk volume is not abundant, it is easier to express by hand than with a breast pump. When milk volume increases, many women find hand expression the easiest way as well.



HAND EXPRESSION IS HELPFUL

- To provide breast milk to a baby that will not latch.
- To relieve breast fullness or engorgement.
- To provide drops of milk to wake a sleepy baby or coax a baby to the breast that is having difficulties latching.
- To help relieve blocked ducts.
- To provide breast milk to rub on sore nipples.
- To soften full breasts to make it easier for baby to latch.
- To increase milk supply by completely draining the breast after a baby finishes nursing or after pumping.

REMEMBER...

- With a newborn it is important to nurse or express milk at least 8 to 12 times per day to maintain a supply.
- Hand expression gets easier with practice and should not hurt.
- The trick to hand expression is discovering where to position your fingers. Experiment until you find what works best for you.
- If you are expressing to increase milk supply, switch back and forth between both breasts. Expressing each side 2 to 3 times may increase milk supply over time.
- If you have concerns about your milk supply, please contact a lactation consultant or breastfeeding expert for assistance.

MISSOURI DEPARTMENT OF HEALTH AND SENIOR SERVICES
WIC and Nutrition Services

P.O. Box 570 Jefferson City, MO 65102-0570 573-751-6204
AN EQUAL OPPORTUNITY / AFFIRMATIVE ACTION EMPLOYER

When Hand Expression is Helpful

TIPS TO HELP MILK FLOW

- Relax and think about your baby.
- If your baby is not with you, look at a picture of your baby.
- Use warm compresses on your breast.
- Practice in the shower with hot water running on your breast – this works well if trying to soften engorged breasts when your baby is 3 to 5 days old.

HOW TO EXPRESS MILK

- Wash your hands and have a clean wide mouth container to catch the milk.
- To begin, gently massage your breasts for a short time to encourage the let-down or flow of milk.
- Position your thumb on one side of your breast and 2 to 3 fingers on the other side; about 1 to 2 inches back from the nipple.
- Press your breast gently inward towards the wall of your chest.
- Compress your thumb and fingers together.
- Release and repeat; catch drops of milk in the container as they appear.
- When milk flow stops rotate thumb and fingers to another position around the nipple and repeat the process until the breast is drained.
- Express milk the same way from the other breast.
- When done expressing, pour breast milk in a clean bottle or storage container.

health.mo.gov/living/families/wic